



## Review

## Keep-ING balance: Tumor suppression by epigenetic regulation

Gesche Tallen<sup>a,1</sup>, Karl Riabowol<sup>b,\*,2</sup><sup>a</sup> Dept. of Pediatric Oncology/Hematology, Charité-Medical School Berlin, Germany<sup>b</sup> Department of Biochemistry & Molecular Biology, Faculty of Medicine, University of Calgary, Canada

## ARTICLE INFO

## Article history:

Received 14 February 2014

Accepted 6 March 2014

Available online 14 March 2014

Edited by Shairaz Baksh, Giovanni Blandino and Wilhelm Just

## Keywords:

ING-protein

Tumor suppression

Epigenetic regulation

Epigenetic cancer therapy

## ABSTRACT

**Cancer cells accumulate genetic and epigenetic changes that alter gene expression to drive tumorigenesis. Epigenetic silencing of tumor suppressor, cell cycle, differentiation and DNA repair genes contributes to neoplastic transformation.**

**The ING (inhibitor of growth) proteins (ING1–ING5) have emerged as a versatile family of growth regulators, phospholipid effectors, histone mark sensors and core components of HDAC1/2 – and several HAT chromatin-modifying complexes. This review will describe the characteristic pathways by which ING family proteins differentially affect the Hallmarks of Cancer and highlight the various epigenetic mechanisms by which they regulate gene expression. Finally, we will discuss their potentials as biomarkers and therapeutic targets in epigenetic treatment strategies.**

© 2014 Federation of European Biochemical Societies. Published by Elsevier B.V. All rights reserved.

### 1. Out of balance: epigenetics in cancer

Cancer has been described in terms of ten specific “hallmarks” that are acquired in different combinations by cells undergoing the multistep process of carcinogenesis [1]. These include mutations in tumor suppressor genes and oncogenes, which have stood the test of time as integral culprits of neoplastic transformation. Oncogenesis may also be understood as a cascade of clonal expansions triggered by random attainment of an enabling mutant genotype [2,3]. Continuously increasing evidence indicates that epigenetic alterations also participate with genetic abnormalities, contributing to the hallmarks that drive the process of tumorigenesis [2,4].

The term “epigenetics” was originally coined in the middle of the twentieth century by Conrad Waddington (1905–1975), a

British developmental biologist. He linked the two fields of epigenesis and genetics to describe “the causal interactions between genes and their products, which bring the phenotype into being” [5–7]. Despite decades of debate, the definitions of “epigenetics” or “epigenetic” remain somewhat controversial [8]. Most commonly, the term is used to describe reversible, chromatin-based events that regulate DNA-templated processes without altering DNA primary sequence [9]. This definition will be used in this article.

The myriad characteristics of a cellular phenotype that are not due to primary DNA sequence alterations evolve by stable maintenance during differentiation, allowing cells to have distinguishable identities while harboring identical genetic information. This means that gene expression patterns are unique, heritable and mediated by chromatin modifications. One major level of chromatin regulation is at the level of the nucleosome, a simple repeating unit of chromatin that contains about 146 bp of DNA wrapped around an octamer of basic histone proteins. Nucleosomal proteins are subject to modification by phosphorylation, ubiquitination, SUMOylation, methylation, citrullination and acetylation, most of which affect histone turnover and/or nucleosome stability. Along with DNA methylation, histone modification has emerged as a promising target in cancer treatment. Acetylation induced by histone acetyltransferases (HAT) is associated with high transcriptional activity, while histone hypoacetylation induced by histone deacetylases (HDAC) usually results in gene silencing. The epigenetic mechanism that is arguably the best understood currently is the transcriptional repression or activation of a

**Abbreviations:** bp, base pair; CDK, cyclin-dependent kinase; DNA, deoxyribonucleic acid; DNMT, DNA methyltransferase; H, histone; HAT, histone acetyltransferase; H3K4me3, histone H3 trimethylated on lysine 4; HDAC, histone deacetylase; ING, Inhibitor of growth; mSin3, mammalian SIN3 homolog; MGMT, O6-methylguanine-DNA-methyltransferase-gene; NFκB, nuclear factor kappa B; NuA4-Tip60, nucleosome acetyltransferase of H 4-Tat-interactive protein-60; PCNA, proliferating-cell nuclear antigen; PHD, plant homeodomain; PtdIns(5)P, phosphatidylinositol 5-phosphate; siRNA, small interfering ribonucleic acid; SWI-SNF, switching deficient-sucrose non-fermentable; SUMO, small ubiquitin-like modifier; Rb, Retinoblastoma-gene; TGFβ, transforming growth factor beta

\* Corresponding author. Address: #311 HMRB, 3330 Hospital Dr. NW, Calgary, Alberta T2N 4N1, Canada. Fax: +1 (403) 283 8727.

E-mail address: [karl@ucalgary.ca](mailto:karl@ucalgary.ca) (K. Riabowol).

<sup>1</sup> MD, PhD, Associate Professor of Pediatrics.

<sup>2</sup> PhD, Professor.

growing list of tumor suppressor genes or oncogenes, by altered methylation of DNA in GC-rich promoter regions (CpG islands) of these genes [9–14].

*A physiological epigenetic balance is maintained when some specialized genes determining the phenotype of differentiated cells are permanently turned on, while others are permanently turned off. Loss of this balance can result in a cancer epigenome, where aberrant transcription of growth regulatory genes and dysregulated signaling pathways enable emergence of the Hallmarks of Cancer.*

In fact, epigenetic imbalance is suspected to be the key initiating mechanism in some forms of cancer [15]. For example, the identification of genes that are specifically hypermethylated (silenced), or hypomethylated (transcriptionally activated), spawned new concepts of cancer initiation, progression and therapy [2,9]. Similar to gene mutations that occur frequently in specific cancers, various high-frequency epigenetic mutations have already been discovered in specific genes. To name a few, the tumor suppressor and growth-regulatory genes Von Hippel–Lindau (VHL) in Wilms Tumor, CDKN2A (encoding the tumor suppressors INK4A and ARF), MGMT (encoding the DNA repair enzyme O(6)-methylguanine-DNA-methyltransferase), Rb as well as the “large tumor suppressor gene 1” (LATS1) are hypermethylated in their promoter regions [16–18]. Another mechanism involved in transcriptional and posttranscriptional gene silencing is specific base pairing between microRNAs (miRNAs) and their growth-regulatory targets. miRNAs are vital for normal development and are frequently compromised in diseases such as cancer [19]. Finally, aberrant epigenetic status in cancers is also defined by posttranslational local and global chromatin modifications [2]. Therefore, studying epigenetics in cancer should not only focus on the discovery of growth-regulatory genes affected, but also on the recognition of the corresponding culprits: chromatin modifying non-histone proteins, that are dysregulated in cancer cells, such as the ING family of proteins that is the major subject of this review.

*In addition to screening cancer-cell genomes and epigenomes for genetic and epigenetic alterations of tumor suppressor genes, cancer patients may profit from the identification and analysis of chromatin-regulatory proteins, since they represent potential biomarkers and targets for epigenetic-based cancer therapies.*

## 2. Almost two decades of learnING: from candidate tumor suppressors to a PHD in epigenetics

The ING (INhibitor of Growth) family of genes and proteins (ING1–ING5) was originally identified in 1996, using subtractive hybridization between cDNAs from a normal mammary epithelial cell strain and several transformed breast cancer cell-lines, followed by an *in vivo* functional screen for tumourigenesis [20]. INGs are evolutionarily well-conserved proteins that localize primarily in the nucleus [21]. Abnormal nuclear morphology is a frequently noted change in cancer cells ever since, more than a century ago, Theodor Boveri described the first genetic alteration in cancer: abnormal chromatin [22,23]. The INGs are multidomain proteins that share motifs targeting them to the nucleus and to different chromatin domains, thereby enabling them to exert their growth-regulatory functions.

### 2.1. All PHDs: on the histone mark, (de)acetylate, go!

All ING proteins share a highly conserved plant homeodomain (PHD) finger (Fig. 1) [21,24,25]. The PHD selectively binds to the lysine 4 residue of histone H3 with affinity increasing with methylation state such that H3K4me3 > H3K4me2 > H3K4me. The H3K4me3 mark is preferentially located at promoters and downstream of transcription start sites [26–34]. It is bound by all ING-PHDs with strong but biologically relevant dissociation constants (*K*<sub>ds</sub> of <10 μM), and competes for binding with a small number of other PHD-containing proteins. Interaction of INGs with H3K4me3, partially by recruitment of the growth arrest DNA damage protein 45a (Gadd 45a), directs the acetylation and methylation status of histone tails, thereby stimulating or silencing growth-inhibitory or pro-proliferative target gene promoters, in multiple species [29,35–39]. Since all of the ING proteins are stoichiometric members of HAT (ING3–ING5) or HDAC (ING1, ING2) complexes, once they bind the H3K4me3 mark through their PHD, they can direct HAT or HDAC activity to the immediate vicinity of the mark to affect chromatin structure.


*Through their conserved PHD, all ING family proteins act as H3K4 methylation-sensitive sensors, thereby transcriptionally regulating target genes. Therefore, they are considered as “readers” of the epigenetic code.*

### 2.2. Nuclear architects sharING motifs

As shown in Fig. 1, ING proteins all encode one or more nuclear localization signals (NLS) that are needed for efficient nuclear targeting [40]. Several of the ING proteins also contain short, highly basic nucleolar translocation sequences (NTS) within the NLS, which translocate them to the nucleoli in response to DNA damage. Mutation of the NTS results in the loss of stress-induced nucleolar targeting and subsequently reduced levels of apoptosis [41].

Many studies to date reporting on the impact of nuclear structure on gene regulation have been performed using developmental systems or aging models such as the Hutchinson–Gilford Progeria syndrome (HGPS), that has a clear link to disruption of nuclear morphology and gene expression by mutation of the gene encoding lamin A. HGPS is a rare developmental autosomal dominant condition characterized by accelerated aging that begins in early childhood and leads to death at a mean age of 13 years [42]. ING1 is one of the genes that is transcriptionally repressed in HGPS and all ING proteins contain a sequence, the lamin interaction domain (LID), that is unique in the entire human proteome. The LID interacts specifically with lamin A and seven other proteins, with lamin A being the highest scoring hit [43]. The interaction between the LID and lamin A is believed to help tether ING(1) in the nucleus, thereby localizing its functions as an epigenetic modifier. Considering that HGPS cells feature aberrant chromatin structure and that ING1 does not interact with the mutant form of lamin A called progerin [43], this raises the possibility that ING proteins contribute to transduction of the HGPS phenotype through altering the epigenetic status of lamin A mutant cells.

Like aging, cancer is associated with global misregulation of gene expression [44–46], suggesting that both aging and cancer are in some aspects, developmental diseases. For example, loss of lamin A due to promoter methylation of the *lamin A/C* gene has been reported for hematologic malignancies [47,48]. While normal B and T cells express lamin A and ING(1), leukemia and lymphoma cells do not, nor do some normal progenitor cells or early B/T cell intermediaries [46,49]. ING2 is overexpressed in colon cancer [50], whereas loss of lamin A is common and a risk factor for



**Fig. 1.** Nuclear architects with shared and unique domains and motifs: structure of ING family proteins. ING family proteins contain a well conserved plant homeodomain (PHD) and lamin interaction domains (LID) through which they function as H3K4me3 sensors, and bind to Lamin A, respectively. A nuclear localization sequence (NLS) that targets them to the nucleus via karyopherin binding also contains basic nucleolar targeting sequences (NTS, black boxes) but these have only been demonstrated to be active in ING1 and so are indicated in gray in other ING members. The NLS region has also been reported to bind p53. The polybasic region (PBR) binds both bioactive phospholipids (PIs) and ubiquitin (Ub) while ARF and SAP30 have been shown to interact with the amino-terminal half of ING1. Proliferating cell nuclear antigen (PCNA) binds specifically to ING1b via the PCNA-interacting protein (PIP) motif, regulating apoptosis and DNA repair in response to DNA damage. The functions of the partial bromodomain (PBD) still have to be characterized. Two phosphorylation sites located on serines 126 and 199 (S126, S199) have been reported to regulate the half-life of ING1 and alter subcellular localization, respectively. *PHD details (rectangular box):* Amino acid sequences of ING PHD zinc fingers that specifically bind H3K4me3 contain Cys and His residues spaced to coordinate two zinc ions (purple rectangles). The enabling Cys and His residues form an aromatic cage (gray rectangles), while other residues contribute specific hydrogen bonds required for recognizing the H3K4me3 histone mark.

relapse in this cancer type [51]. Thus, if the ING proteins function as effectors localized by lamin A, colon cancer cells may upregulate ING2 to compensate for lamin A deficiency.

*Aging and cancer share many common features of a perturbed nuclear architecture and aberrant gene expression that may be partially due to dysregulated interactions between lamin A and ING proteins.*

### 2.3. Unique motifs for linING tumor suppression to transcriptional and post-translational regulation

ING1 also contains a *partial bromodomain (PBD)* [21] (Fig. 1). This motif can interact with SAP30 of the Sin3-HDAC1 and HDAC2 complexes [52], thereby regulating transcription [32,53]. Furthermore, a *polybasic region (PBR)* adjacent to the PHD, is unique for ING1 and ING2 (Fig. 1). Within the human proteome, the ING2 protein shows the strongest affinity for bioactive phosphoinositides such as phosphatidylinositol 5-phosphate (PtdIns(5)P) in the nucleus. This interaction regulates the growth inhibitory effects of ING2 in a p53-sensitive manner [54,55] and is of biological significance for the following reasons: phospholipids that specifically bind to ING1 and ING2 are stress-inducible and highly bioactive. This may explain why ING1 and ING2 proteins targeting HDAC complexes to chromatin are strongly induced in response to stress. Additionally, the PBR overlaps with a *ubiquitin-interaction motif (UIM)* in the C-terminal region of the ING1b isoform [56] (Fig. 1). Ubiquitin and phosphorylated lipid species compete for binding to this site, thereby linking bioactive lipid signaling to ubiquitin-mediated proteasomal degradation [56]. The PBR/UIM of ING1b also interacts with mono-ubiquitinated p53. As a result, the p53 tumor suppressor is stabilized, most likely by blocking polyubiquitination, through targeting deubiquitination by the herpesvirus-associated ubiquitin-specific protease HAUSP [56].

With the exception of ING1, all ING proteins also have a *leucine zipper-like region (LZL)* [21] (Fig. 1) that can help regulate nucleotide-excision-repair-associated functions of ING2, the DNA damage response and cell differentiation [57]. The LZL of ING2 also interacts with components of HDAC1 complexes and it is a critical domain in influencing muscle differentiation by ING2 [57].

The *proliferating-cell nuclear antigen (PCNA)-interacting protein (PIP)* is a domain that is unique to ING1b (Fig. 1). ING1b interacts with PCNA in a UV-inducible manner to initiate DNA repair and/or apoptosis as a stress response [58]. ING1b also contributes to E3 ubiquitin ligase Rad18-mediated PCNA mono-ubiquitination. Mono-ubiquitinated PCNA associates with DNA polymerase  $\eta$  (Pol $\eta$ ) to promote proper lesion bypass and error-free DNA replication [59].

Last but not least, recently published results indicate that another domain unique for ING1 and ING2 exists: a *SUMOylation-motif (PDSM)*, which provides interaction of ING proteins with the E3-type small ubiquitin-like modifier (SUMO) ligase PIAS4 that has implications in the DNA damage response [60].

*Due to their structural characteristics, ING family proteins are nuclear proteins with the ability to act as:*

- histone mark sensors connecting cellular responses to genotoxic stress,
- stoichiometric members of HAT and HDAC complexes,
- growth regulators.

### 2.4. Family meetING

ING family proteins were originally characterized as tumor suppressors based on observations obtained in the era before the development of siRNA technology [61]. Over the last decade, their definition has broadened due to results from multiple siRNA studies and in vivo knockout models. These revealed that ING family proteins differentially control cell growth as epigenetic regulators in different biological contexts.

### 2.5. ING1 – ImprovING with age

The founding member of the ING family, *ING1*, is localized on chromosome 13q34 [21]. The major ING1-isoforms, ING1a and ING1b, are widely expressed in normal and cancer cells [62,63]. They differ in their N-termini (Fig. 1) and growth-regulatory effects, with ING1a affecting cell senescence [64,65] and ING1b contributing to apoptosis [54,58,66–69]. Over almost two decades of study, ING1 was initially identified as a type-II tumor suppressor, then also as an inducer of replicative senescence and more recently as a transcriptional regulator.

#### 2.5.1. ING1 in human cancers

*ING1* gene and/or protein expression is down-regulated in multiple malignancies. These include childhood acute lymphoblastic leukemia [49], neuroblastoma [70], melanoma [71–74], lung [75–78], ovarian cancer [75,78], malignant glioma [79,80], gastric [81], colorectal [82], head and neck squamous cell carcinoma (HNSCC) [83,84], pancreatic [75,85], prostate [75], and breast cancer [75,86,87]. ING1 has also been identified as a breast cancer antigen by serological analysis of recombinant tumor cDNA expression libraries [88]. ING1-overexpression correlated with inhibition of metastasis and reduced tumor-induced mortality of breast cancer *in vivo* [89]. It was also associated with poor prognosis in neuroblastoma [63] and bladder cancer [90]. However, most of these reports do not include information on ING1's subcellular localization. Since ING proteins, like p53 [91], have both nuclear and cytoplasmic functions, localization of ING1 should be confirmed before establishing correlations between expression levels and clinical data.

#### 2.5.2. Mechanisms of ING1 regulation

*Ing1*-deficient mice are characterized by reduced body size, as well as earlier onset and higher incidence of lymphomas [66,67,92–94]. Moreover, high levels of ING1 were observed in regions of mouse embryos, which are known for increased apoptosis during embryogenesis [95]. Thus, *ING1* downregulation in human cancers may simply reflect a less differentiated phenotype. Alternatively, ING1 loss could itself contribute to the neoplastic process. *ING1*-downregulation was not due to mutation, but frequently correlated with reduced ING1 protein levels [70–87]. These observations are indicative of epigenetic mechanisms regulating *ING1* transcription. *ING1* is very GC-rich in four regions that make it a prime target for CpG-island methylation [96,97]. Indeed, promoter methylation has been reported for *ING1* in ovarian cancers [75] and chronic lymphoblastic leukemia [98]. Abnormal methylation patterns in other genes can also result in *ING1* downregulation, as it has been shown for murine adult brain cells [99]. Intriguingly, the microRNA miR-622 is, like ING1 [81], downregulated and involved in invasion and metastasis of gastric cancer [100]. Moreover, up-regulation of miR-622 repressed *ING1*-transcription in this cancer type [100]. Post-translationally, various growth-inhibitory effects of ING1 are regulated by phosphorylation at Ser126 [101,102]. Also, phospho-serine-mediated 14-3-3 binding and nuclear export regulate the ability of ING1b to transcriptionally regulate the p21



cyclin-dependent kinase inhibitor and subsequent induction of apoptosis in response to DNA damage [103]. Moreover, the proto-oncogene Src, a non-receptor tyrosine kinase with an important role in growth factor signal transduction, phosphorylates ING1b, relocalizes it to the cytoplasm, thereby regulating its pro-apoptotic abilities [104]. Overall, valid interpretation of *ING1* downregulation by cancers requires more studies of *ING1* promoter methylation status, of subcellular *ING1* protein localization as well as co-expression analyses of other growth-regulators, including additional miRNAs, of which *ING1* may be a *bonafide* target.

*ING1* is downregulated but not mutated in various human cancers. *ING1* expression is regulated by epigenetic mechanisms, such as promoter methylation, miR-622 and by PTMs such as stress-inducible phosphorylation by Src.

### 2.5.3. Affecting cancer hallmarks by epigenetic regulation


Overexpression of human *ING1* causes cell-cycle arrest in the  $G_0/G_1$ -phase, whereas suppression leads to loss of cellular growth control and immortalisation [20]. While *ING1b* is mainly involved in regulating apoptosis, cellular stress, hormonal responses as well as inhibition of tumor angiogenesis [105–107], *ING1a* inhibits growth by inducing replicative senescence, partially through inhibiting endocytosis via the Rb-tumor-suppressor pathway [64,65]. *ING1b* can efficiently induce apoptosis through interaction

with other tumor suppressors like p53 [68,108,109] and with members of the intrinsic apoptotic pathway such as BCL-2-associated X protein (BAX) [69]. Moreover, *Ing1* inhibits proliferation of mature B cells in mice [110]. In *Ing1* knockout mice, however, loss of p37*Ing1* induces Bax expression and promotes DNA damage-induced apoptosis in a p53-independent manner [110]. *Ing1*-loss also increases sensitivity to ionizing radiation and is associated with earlier onset and higher incidence of lymphomas in mice [110]. This may be due to the involvement of *Ing1* in regulating various DNA repair mechanisms in response to stress [111]. These observations suggest both tumor suppressive and oncogenic effects of *ING1*, possibly depending on the species and biological context.

### Major Hallmarks of Cancer affected by *ING1* (Fig. 2):

- dysregulated growth suppression,
- replicative immortality,
- invasion and metastasis,
- resistance to cell death,
- aberrant proliferative signaling,
- pathological angiogenesis.

This potentially dual function of *ING1* may be due to its function as an epigenetic regulator. *ING1* can both activate and repress target genes, since it is an H3K4me3 sensor that



**Fig. 2.** Left panel: hallmarks of cancer affected by *ING*-HDAC- and HAT-complexes. White shapes and symbols clockwise (according to Hanahan and Weinberg 2011) [1]: arrow: aberrant proliferative signaling, “No”-symbol: dysregulated growth suppression, Y: immune destruction, ∞: replicative immortality, flame: tumor-promoting inflammation, triangles: invasion & metastasis, pipe: pathological angiogenesis, DNA: genome instability & mutation, cross: resistance to cell death, “O<sub>2</sub>”: deregulated cellular energetics, red “stop”-sign: inhibited by *ING* activity, question-mark: possibly inhibited by *ING*(5) activity. Abbreviations: HAT – histone acetyl transferase, HBO1 – HAT of the MYST family, HDAC1/2 – histone deacetylase 1/2, *ING* (1–5) – Inhibitor of growth (1–5), JADE – zinc finger protein associating with MYST-HATs, mSin3A – core component of large multiprotein HDAC-associated corepressor complex, Tip60-MYST – HAT of the MYST family. Subunits/domains not explained in the text: BRMS1 – breast cancer metastasis suppressor 1, BRPF1/2/3 – PHD proteins bridging MOZ and MORF in MYST-HAT-complexes, EPC1/2 – ectopically parting cells 1/2, hEaf6 – subunit of MYST-HAT-complexes. For detailed information on proteins and signaling pathways associated with the tumor suppressive functions of *ING* proteins: see main text. Right panel: Stress signaling and chromatin modification by *ING* family proteins. (1) **Activation:** upon cellular stress, p38MAPK is activated and then translocated to the nucleus where it phosphorylates PIP4Kβ, thereby inhibiting its kinase activity and increasing nuclear PIP levels. PIP subsequently binds to *ING1* and *ING2*. Recognition of the histone mark H3K4me3 by *ING* proteins results in formation of *ING*-HDAC or -HAT complexes. (2) **Interpretation:** *ING* proteins specifically bind to H3K4me3 or H3K14me3 recruiting the assigned chromatin remodeling complexes to specific chromatin loci and target gene promoters (details in main text). This results in various distinct histone-PTMs, such as acetylation, deacetylation and methylation as well as subsequent activation (gene “on”) or silencing (gene “off”) of gene transcription affecting the Hallmarks of Cancer [1]. (3) **Migration:** after realization of histone modifications by *ING*-HDAC and HAT-complexes, *ING* proteins are removed and/or inactivated by different PTMs [24]. These include phosphorylation-dependent binding to 14-3-3 proteins leading to cytoplasmic translocation, whereas dephosphorylation releases *ING* from 14-3-3, thereby promoting *ING* binding to karyopherin proteins κ $\alpha$  or κ $\beta$  and subsequent re-localization of *ING* to the nucleus. Abbreviations and symbols: ac – acetyl group, me3 – 3-methyl group, HAT – histone acetyl transferase, HDAC – histone deacetylase, *ING* – Inhibitor of growth, K4, K14 – core histone subunits, PI – phosphatidylinositol, PIP4Kβ – phosphatidylinositol-5-P 4-kinase type 2 beta, PIP – phosphatidylinositol monophosphate, PTM – post-translational modification, RNA-PolII – RNA polymerase II; yellow pentagons: phosphorylated sites.

recruits mSin3A/HDAC1/2 [52] (Fig. 2). It also cooperates with numerous proteins that are important for different epigenetic modulations, including the Ras, p300 and p16(INK4a) as well as DNA methyltransferase 1 associated protein (DMAP1) [112–116]. Gadd45a requires specific ING1 binding to H3K4me3 for DNA demethylation [38], thereby linking histone methylation to DNA demethylation. Moreover, ING1 is a transcriptional regulator of multiple miRNAs: ING1b overexpression induces miR-203, miR-375, miR-449b and miR-200c, thereby downregulating Src and CDK6, to name a few, and promoting growth inhibition effects [117]. In mouse primary embryonic fibroblasts, Ing1 regulates the microRNA regulator protein Dgcr8, a protein involved in the early steps of microRNA synthesis [118]. These findings establish a new link between tumor suppression and microRNA biogenesis.

#### ING1 regulates target gene expression by

- recruitment of protein complexes with chromatin-modifying activity to specific (H3K4me3-marked) promoters,
- regulating Gadd45a-mediated DNA demethylation,
- regulating miRNA expression signatures and biogenesis.

### 2.6. ING2: The Renaissance Protein

ING2 has high amino acid sequence homology to ING1 [21,119,120] (Fig. 1). Two major splicing variants have been identified for ING2 (ING2a, ING2b) [121], with the original “ING2” now called ING2a, when required. ING2b lacks the N-terminus that determines major functions of ING2a [122]. ING2 is characterized by its versatility: it is another histone mark sensor that can act as a DNA damage responder and phospholipid signaling effector [123]. In addition to its growth-regulatory abilities, ING2 possesses essential developmental functions, particularly in spermatogenesis [93], axon growth, neuronal and muscle differentiation that are mediated by p53- and chromatin-mediated mechanisms [124], as well as by SnoN-induced transcriptional regulation through ING2 [125].

#### 2.6.1. ING2 in human cancers

Expression of ING2 is upregulated in colorectal cancer [50,120], in Burkitt lymphoma and cervical cancer according to the Oncomine Database (<http://www.oncomine.org/main/login.jsp>). Decreased ING2 expression or loss of heterozygosity have been found in various types of malignancies as well. These include hepatocellular carcinoma (HCC) [126,127], HNSCC [128] and lung cancer [75,129,130]. The subsequent conclusion that ING2 is generally downregulated in these cancers still needs to be confirmed for the following reasons: most of the associated studies were carried out using either a relatively small number of cell-lines [75], or revealed reduced expression of ING2 at the mRNA level, but no significant changes at the protein level [126]. Despite this, and although subcellular localization of ING2 was cytoplasmatic, even in normal control tissues, ING2 protein expression correlated with unfavorable prognosis [126]. However, ING2 is a nuclear protein [122]. This, as well as additional controls to verify antibody staining specificity should be considered before using expression data in Kaplan–Meier analyses.

#### 2.6.2. Mechanisms of ING2 regulation

Aberrant ING2 levels in cancers appear not to be due to mutations. Instead, direct binding of the p53 tumor suppressor to the promoter region of ING2 results in ING2 downregulation [131]. Considering that about 50% of sporadic cancers are

characterized by TP53 mutations [132], the upregulation observed for ING2 in colorectal cancers might be, as outlined above, a feedback mechanism due to lamin A loss [51] and/or p53-deficiency. On the other hand, Nutlin 3a, an MDM2 inhibitor that activates p53, is known to downregulate endogenous ING2 levels through binding to two p53 binding sites on the ING2 promoter. This interaction causes senescence in normal cells [131] and could be a mechanism by which ING2 levels are being reduced in the remaining other 50% of malignancies with wild-type p53. Hence, interpretation of ING2 levels in tumors should also include their p53-status. ING2-upregulation may also be a result of direct binding of the transcription factor nuclear factor kappa B (NF-κB) to the ING2 promoter [133]. NF-κB is antiapoptotic and, like ING2, is overexpressed in colorectal cancer [133]. ING2 protein levels are also regulated by PTMs, and in particular ubiquitin- and SUMO-mediated mechanisms. ING2 interacts with ubiquitin ligase Smad ubiquitination regulatory factor 1 (Smurf1) that targets ING2 for poly-ubiquitination and proteasomal degradation [134], and is regulated by SUMOylation, affecting Sin3A activity [135].

*ING2 is differentially expressed in different cancers, transcriptionally repressed by Nutlin3a-mediated p53-activity and activated by NF-κB.*

*ING2 proteins are post-translationally modified by Smurf-mediated ubiquitin-dependent proteasomal degradation.*

#### 2.6.3. Affecting Cancer Hallmarks by epigenetic regulation

ING2 is a damage-inducible gene. The ING2a isoform negatively regulates cell proliferation through p53 activation via acetylation mediated by the HAT p300 [136] and by regulating p21 independently of p53 [137]. ING2 also mediates transforming growth factor-β (TGF-β)-dependent responses in epithelial cells. Considering that TGF-β has growth-inhibitory functions in normal cells and oncogenic functions in invasive metastatic malignancies, the signals from TGF-β mediated by ING2 appear to differ in normal and cancer tissue [138,139]. Moreover, ING2a is required for the initial DNA damage sensing and chromatin regulation in the nucleotide excision repair process via recruitment of the damage-recognition Xeroderma pigmentosum group A-complementing (XPA) protein to the lesion site after DNA damage by UV [140]. In contrast, ING2b alone appears not to have any growth-regulatory effects [122]. Thus, the two isoforms may have compensatory roles that protect cells from cell cycle arrest and apoptosis, as only loss of both can induce these biological functions, particularly in p53-deficient cells. These observations imply a role for ING2 proteins in the development of resistance to conventional cancer treatments, such as DNA damaging agents. Moreover, differential regulation of the two ING2 isoforms may be essential for maintenance of irreversible cellular senescence seen following the induction of high levels of DNA damage.

#### Major Hallmarks of Cancer affected by ING2 (Fig. 2):

- dysregulated growth suppression,
- replicative immortality,
- invasion and metastasis,
- resistance to cell death,
- aberrant proliferative signaling.

Loss of Ing2 in Ing2-knockout mice resulted in male sterility and increased incidence of soft tissue sarcomas [124]. In contrast, ING2 loss in human gastric cancer cells induced G<sub>0</sub>/G<sub>1</sub> arrest and

reduced their invasive behavior [141]. ING2 appears to have, similar to ING1, differential effects in mice and men, in normal and in cancer cells – a role that is consistent with an epigenetic regulator. Moreover, the PHD finger of ING2 integrates phosphoinositide and chromatin signaling networks to prevent unchecked cell growth: Upon DNA damage, ING2 binds to H3K4me3 [142,143] mediated by nuclear PtdIns(5)Ps, thereby recruiting mSin3a/HDAC1 [144] (Fig. 2). Subsequent promoter binding of the ING2-SIN3A-HDAC1-complex is regulated by SUMOylation of ING2 [135]. To date, more than 200 gene promoters have been identified that are repressed or activated by ING2 in response to genotoxic stress [146]. For example, ING2 silences the candidate proto-oncogene *CIP2A* [KIAA1524] [144], whereas transcription of *matrix metalloproteinase 13* (*MMP13*) can be activated or repressed by ING2, thereby regulating the invasion ability of cancers [133,145]. ING2 also binds to the p21 promoter to control the G<sub>1</sub>/S checkpoint [139]. It regulates cell cycle progression by blocking the function of RBP1: RBP1 allows recruitment of the mSin3A-HDAC complex by Rb pocket proteins to induce cell cycle arrest through repression of E2F-dependent transcription and DNA replication origins [146]. RBP1-associated mSin3A-HDAC1 activity is inhibited by SIRT1, which is recruited by ING1 and ING2 proteins [147]. In addition to regulating gene transcription by HDAC- and HAT crosstalk, ING2 collaborates with the tumor suppressive SWI-SNF-BRG1-complex. This is similar to ING1 interacting with the corepressor Alien in chromatin remodeling and both recruit histone methyltransferase (HMT) activity to methylate histone H3 [53,138,148,149].

#### ING2 regulates target gene expression by

- stress-inducible, phosphoinositide-mediated recognition of H3K4me3, subsequent recruitment of the mSin3A/HDAC1 complex and binding of target gene promoters,
- chromatin remodeling via association with SWI/SNF/BRG1,
- methylation of histone H3 through HMT recruitment.

### 2.7. ING3 likes MYSTerious HATs

ING3 was identified through bioinformatic analyses of the human genome [62,150]. Its amino acid sequence is the most distinctive among the five ING proteins evolutionarily (Fig. 1) [21]. ING3 modulates cell growth and p53-mediated transcription [150,151]. Although ING3 is ubiquitously expressed in mammalian tissues, it has been reported that it is more highly expressed in oocytes from mice, rhesus monkeys, and humans [152]. Moreover, ING3 has the highest molecular weight in the family due to long unique, and still uncharacterized regions. These imply unique roles for this family member, such as linking developmental functions to chromatin structure, as recently found in germ and blood cells [152–155].

#### 2.7.1. ING3 in human cancer

Various studies of cancer samples and cell lines analyzed for ING3 mutation and expression have reported that the ING3 gene was silenced. These include HNSCC [156,157], HCC [158,159] and ovarian cancer [75].

#### 2.7.2. Mechanisms of ING3 regulation

ING3 is a miR-21 target gene with a role in monocyte differentiation to dendritic cells [155]. Additionally, ING3 represents a novel transcription target in the serine/threonine kinase RSK2-cAMP-response element-binding protein (CREB) pathway: RSK2 signaling through CREB downregulates ING3 to protect HNSCC cells from the apoptotic “anoikis” process, thereby contributing

to cancer cell invasion and tumor metastasis [160]. On the protein level, ING3 is acetylated on several lysine residues that are all located in the unique region of the protein [161]. Thus, ING3 might be distinctively regulated by acetylation. Like ING1 and ING2, ING3 is also ubiquitinated leading to proteasomal degradation [102,162].

*ING3 is downregulated in human cancers and transcriptionally regulated by miR-21 and RSK2 signaling through CREB.*

*ING3 is post-translationally modified by acetylation and ubiquitination at Lys residues within its unique regions.*

#### 2.7.3. Affecting Cancer Hallmarks by epigenetic regulation

Numerous reports consider ING3 as another tumor suppressor in the family, since it promotes apoptosis in human cancer cells [148,150]. However, these observations mostly originate from unphysiological overexpression experiments and may require confirmation by knockdown models. In *Caenorhabditis elegans*, however, *ing-3* downregulation by siRNA resulted in strong suppression of stress-induced programmed cell death in germline cells and in the developing embryo [153].

#### Hallmarks of Cancer potentially affected by ING3 (Fig. 2).

- resistance to cell death,
- aberrant proliferative signaling.

The epigenetic activities of ING3 support its potential role as a growth inhibitor in humans: In addition to binding to H3K4me3 similarly to the other ING family members [148], ING3 is a member of the nucleosome acetyltransferase of H4-Tat-interactive protein-60 (NuA4-Tip60)-MYST-HAT complex that acetylates the N-terminal tails of histones H4 and H2A [142,143,148,163] (Fig. 2). The MYST proteins are part of large conserved multisubunit HAT complexes that have diverse roles in gene expression, carcinogenesis, tumor progression, DNA replication and DNA repair [164]. Because the NuA4-Tip60 complex is a MYST-HAT complex that acetylates histones H4 and H2A [165], ING3 may bind to trimethylated lysines of these histones as well. Since histone acetylation and H3K4me3 recognition associate with transcriptional activation, ING3 may be a transcriptional activator only, unlike ING1 and ING2, which can be involved in both transcriptional activation and suppression. Genes regulated by ING3 have yet to be reported.

*ING3 is a histone mark (H3K4me3) sensor and member of the NuA4-Tip60-MYST-ING-HAT complex that may activate target genes by histone H2A and H4 acetylation.*

### 2.8. ING4 – that's not old HAT but a HAT-trick!

ING4 was identified by computational homology search [62]. Several alternative transcripts of ING4 have been found so far [166,167] (Fig. 1). One of them (ING4\_vs4) was initially designated an ING4 mutation unique for cancer cells until it was recognized as a splice variant [168]. All ING4 isoforms are ubiquitously expressed in various tissues [166]. The roles of each variant still have to be elucidated, however, designing a specific siRNA against each variant remains challenging. ING4\_v2, ING4\_v3, and ING4\_v4 have 1, 3, and 9 amino acids deleted in the NLS, which are important for



p53 binding [169]. Reports on the effects of small deletions in the NLS on subcellular localization are controversial [166,165,170]. Thus, ING4 isoforms may distribute differentially in different cell types and different biological contexts, and interact with different binding partners to transduce different activities of ING4 during carcinogenesis.

### 2.8.1. ING4 in human cancers

ING4 is mainly lost or downregulated at the RNA level in human cancers. These include astrocytic tumors of the central nervous system [171,172], breast cancer [75,168,173–175], malignancies of the gastrointestinal tract [176–178], HNSCC [179,180], HCC [181,182], melanoma [75,183], lung [75], ovarian [75] and prostate cancer [75]. Like ING1 [80], ING4 downregulation correlates with higher WHO grades of malignancy in astrocytoma [172], with unfavorable prognosis of breast cancer patients [89,173,184] and with angiogenesis in multiple cancer types [106,184–190] and other diseases [191].

### 2.8.2. Mechanisms of ING4 regulation

ING4 is downregulated by miR-650 or miR-214, respectively, in leukemia, HCC, lung, gastric and pancreatic cancer [186,192–194], thereby possibly contributing to resistance to apoptosis as well as to certain anticancer agents [195]. ING4 protein stability and activity are primarily regulated post-translationally by citrullination on the NLS-motif. ING4 citrullination alters ING4 half-life, reduces its ability to acetylate p53 and promotes ING4 degradation [196].

*ING4 is downregulated in human cancers and transcriptionally regulated by miR-214 and miR-650. ING4 is post-translationally modified by citrullination on the NLS-motif.*

### 2.8.3. Affecting Cancer Hallmarks by epigenetic regulation

In concert with its multiple binding partners such as p53, the HAT p300, HPH-2, which regulates HIF- $\alpha$  stability, the p65 subunit of NF- $\kappa$ B, and liprin  $\alpha$ 1 [61,197], ING4 primarily acts as a tumor suppressor. Exogenous ING4 reduces cell colony formation, decreases S-phase in cycling cells [138,168], and induces cell death in cancer cells by autophagy [197] or p53-mediated apoptosis [198]. *Ing4*-null mice, however, are viable, develop normally and do not form spontaneous tumors [199]. But *in vivo* models also revealed that Ing4 plays a major role in the inflammatory response of mice to bacterial components through its ability to suppress NF- $\kappa$ B activation of select cytokine genes in stimulated macrophages [199]. Collectively, these results suggest a previously unexplored role also for the human counterpart ING4 in negatively regulating NF- $\kappa$ B-mediated innate immunity. Taken together, ING4 appears to particularly target the three cancer hallmarks angiogenesis, immune destruction and tumor promoting inflammation [1](Fig. 2).

#### Major Cancer Hallmarks affected by ING4 (Fig. 2):

- immune destruction,
- tumor promoting inflammation,
- angiogenesis.

In addition, ING4 binds to H3K4me3 [200] to associate with the MYST-HB01-JADE-hEAF6-HAT-complex [148,201] (Fig. 2). This latter complex is responsible for most nucleosomal histone H4 acetylation in eukaryotes, and knockdown experiments indicated

that Ing4-HB01 association is required for cells to progress properly through the S-phase of the cell cycle [165]. Moreover, ING4 bridging HB01 with JADE proteins promotes transcriptional activation of erythroid developmental target genes [202]. Although both the ING3 complex and the ING4 complex include MYST-HATs, these acetylases are different between the complexes: the ING3 complex includes Tip60 and the ING4 complex includes HB01 as a catalytic subunit. The ING4-HB01 complex acetylates histone H4K16, whereas the Tip60-ING3 complex does not acetylate this histone residue [165]. These findings indicate that each ING family protein has a different elaborate role in transcriptional regulation through generation of different “histone code” signals.

*ING4 is a histone mark sensor and a member of the HB01-JADE-hEAF6 MYST-ING HAT-complex, thereby activating target genes by acetylation of histone H4.*

### 2.9. Lift two HATs to ING5!

ING5 was identified by computational homology search and shares high amino acid sequence homology with ING4 (Fig. 1) [62,198] ING5 associates with HAT complexes and, depending on the biological context, may have tumor suppressive or oncogenic abilities (Fig. 2).

#### 2.9.1. ING5 in human cancers

A few lung, pancreatic and ovarian cancer cell lines [75], but also bone marrow samples from a large cohort of patients with acute myeloid leukemia (AML) revealed *ING5* downregulation, hinting at a potential tumor suppressive effect of ING5 in these malignancies [203]. In breast cancer, however, only 2/9 cell lines showed aberrant, i.e. decreased *ING5* gene expression [75]. In the majority of gastric cancer samples, ING5 is downregulated at the RNA-, and upregulated on the protein level [204], which may be due to posttranslational dysregulation. *ING5* is also downregulated in HNSCC, in which it has been identified as a tumor suppressor [205,206]. Moreover, cytoplasmic localization of ING5 was significantly increased in these tumors and inversely correlated with nuclear ING5 levels that predict a well-differentiated status. Nuclear localization of ING5 positively correlated with p21 and p300 expression, and with the apoptotic index, suggesting that subcellular mislocalization of ING5 may modulate the transactivation of target genes, thereby compromising apoptosis and cell cycle arrest.

In contrast to the cases noted above, *ING5* gene expression is upregulated in colon cancer [61], suggesting an oncogenic function of ING5 in these malignancies. However, these findings might also reflect that the subcellular localization of ING5, which was not part of this analysis, is of major relevance for its growth regulatory effects: a subsequent study [207] revealed no aberrant *ING5* gene expression, but upregulated protein levels in more than 70% of patients samples and colon cancer cell lines. Moreover, nuclear ING5 negatively correlated and cytoplasmic ING5 positively correlated with aggressive behavior of the tumors. Thus, both information on expression and subcellular localization of ING5 is required for assessing its role in the biology of colon, and most certainly other cancers. Moreover, expression of REIC [208], a member of the Dickkopf (Dkk) family, might be of relevance when interpreting ING5 expression in colon cancers; REIC/Dkk-3 acts as a tumor suppressor in multiple cancer cell lines by inducing apoptosis through endoplasmic reticulum stress signaling, and significantly negatively correlated with ING5 expression in colon cancers [208]. However, further analysis is needed before establishing whether a functional ING5-Dickkopf interaction exists.



### 2.9.2. Mechanisms of *ING5* regulation

In pancreatic cancer, *ING5* is negatively regulated by miR-196a, upregulation of which is associated with poor prognosis [209]. In mesenchymal stem cells (MSC), *ING5* mRNA and protein levels are regulated by miR-193 to control CDK2-mediated MSC proliferation [210]. *ING5* protein activity is further controlled by methylation [211] and acetylation [161].

*ING5 is differentially expressed in different cancers and transcriptionally regulated by miR-196a. ING5 proteins are post-translationally modified by methylation and acetylation.*

### 2.9.3. Affecting cancer hallmarks by epigenetic regulation

Being a genuine ING, *ING5* controls growth differently in different biological settings: *ING5* induces cell cycle arrest and apoptosis in normal and cancer cells [148,198,203,206,209]. In murine mesenchymal fibroblasts, *Ing5* requires Inhibitor of cyclin A1 (INCA1) to promote Fas-induced apoptosis [207]. *ING5* is also involved in p53-dependent stress-signaling, partially by physically interacting with p53, p300/CBP- and MYST-HATs (Fig. 2), such as HBO1 and Tip60, thereby inducing the activation of p53-downstream effectors [148,198,212]. *ING5* also associates with minichromosome maintenance (MCM) proteins. These play an essential role in DNA replication [148]. Moreover, knockdown of *ING5* completely inhibited DNA synthesis, whereas knockdown of the HAT HBO1 increased cells in S-phase. Thus, the HBO1-JADE-*ING5*-HAT-complex may be involved in carcinogenesis by enhancing DNA replication. In normal cells, *ING5* inhibition by siRNA resulted in increased MSC proliferation [210], and was found to be part of a network of epigenetic modifiers that regulate epidermal stem cell differentiation [213].

#### Major Hallmarks of Cancer affected by *ING5* (Fig. 2):

- dysregulated growth suppression,
- resistance to cell death,
- invasion and metastasis,
- aberrant proliferative signaling.

Like other ING family proteins, *ING5* is a H3K4me3 sensor [142,143,214], thereby most certainly regulating both, gene silencing and activation. *ING5* is also a component of two different HAT complexes [151]. Depending on its catalytic acetyltransferase subunit, i.e. MOZ/MORF or HBO1, it either binds to histone H3 or H4, respectively [148]. As a subunit in the HBO1-MYST-ING-HAT-complex that usually contains BRPF1, *ING5* can disassemble BRPF1 from unmethylated H3K4, thereby switching HBO1-specificity from H3 to H4 tails [215]. This suggests a unique role for *ING5* as a regulator of HAT affinity to specific histone tails.

*ING5 is a histone mark (H3K4me3) sensor and member of the HBO1-JADE-hEAF6- and MOZ/MORF-MYST-ING-HAT-complexes, thereby activating target genes by acetylation of histones H3 and H4.*


### 2.10. Target-ING individual cancer epigenomes: possible implications for the ING family proteins in epigenetic cancer therapy

Current multimodal treatment concepts for cancer patients are limited due to toxic side effects in normal cells and frequent

development of resistance in cancer cells. Newer strategies target well-defined molecular markers, such as overexpressed Her-2 in breast cancer or Bcr-abl fusion in leukemia. They are often effective initially, but fail when subpopulations of resistant cells evolve. Thus, the new paradigm of anti-cancer drug development aims at combating multiple Hallmarks of Cancer simultaneously. In this context, epigenetic-modulating agents harbor a variety of options, since they can (a) reverse stem-cell like behavior and chemoresistance and (b) simultaneously affect multiple signaling pathways that might be dysregulated in a certain cancer type. Moreover, every patient, thus every individual cancer, has a unique histone code, modification of which is reversible, thereby opening a wide therapeutic window and leaving the genetic code unaffected. Hence, epigenetic “reprogramming” of a cancer cell by either reactivating silenced tumor suppressors or by silencing oncogenes may sensitize cancer cells to lower doses of conventional cytotoxic agents or reverse resistance to other targeted approaches. Indeed, anticancer drugs targeting DNA methyltransferases (DNMTs) and HDACs have demonstrated antitumor activity in the clinic. To date, the best characterized and FDA-approved epigenetic modulating antineoplastic agents are DNMT inhibitors (DNMTi) and HDAC inhibitors (HDACi) for treatment of hematological malignancies. Success in solid tumors has been more elusive, possibly due to ineffective delivery, dosing or scheduling [216]. Hence, different strategies for combining a DNMTi such as 5azaC with other chemotherapeutics like DNA damaging agents or an HDACi, are currently under investigation. Moreover, recent reports on the roles of the bromodomain-containing BET family members and methyltransferase EZH2 in driving cell growth and survival have provided further validation of epigenetic regulators as critical drivers of the transformed cancer cell phenotype [217]. Similar to BET proteins, ING proteins are components of the write-read-erase concept that has been linked with the transfer of epigenetic information. Therefore, they represent both potent biomarkers and a plethora of potential novel targets for future epigenetic cancer therapy. A model for their possible implications is presented in Fig. 3.

#### 2.10.1. Target-ING the mSin3A-HDAC-complex: implications for *ING1* and *ING2*

Cell death and caspase 3-mediated apoptosis in breast cancer cell lines significantly increased when combining 5azaC and the HDACi LBH589 with adenoviral delivery of *ING1b* [82]. This effect was not significantly dependent on p53-status [89]. It was strongest in estrogen receptor (ER)-negative breast cancer cells, possibly because *ING1b* can activate transcription of ER $\alpha$  [218,219]. *In vivo*, *ING1* plus 5azaC significantly reduced the sizes of subcutaneously xenografted tumors in SCID mice without significant toxicity or development of resistance [89]. These observations suggest a combination of *ING1*, DNMTi and HDACi as a promising approach for patients with ER-negative breast cancer. In glioblastoma multiforme (GBM), a common and highly malignant brain tumor, downregulation of *ING1* by siRNA resulted in reduced FADD and caspase-3 mediated apoptosis by the HDACi trichostatin A (TSA) [220]. Because neither FADD nor caspase 3 was observed to be a transcriptional target of *ING1* [221,222], *ING1* function may rather be to stabilize or activate members of this apoptotic pathway. Given the fact that major tumor suppressor pathways are already functionally eliminated in most GBM [223], maintaining *ING1* levels in these cells may allow HDACi-induced apoptosis to be of major therapeutic impact. Considering that *ING1* expression is significantly reduced in malignant gliomas [79,80], and that downregulation of *ING1* by siRNA also sensitizes p53-deficient malignant glioma cells to cisplatin-induced cell death [224], *ING1* levels in GBM cells might influence the choice of chemotherapy. This strategy may also apply to patients with osteosarcoma, since *ING1b*



**Fig. 3.** Balanc-ING the cancer epigenome: ING family proteins as potent therapeutic targets for targeting the Hallmarks of Cancer. *Abbreviations and symbols:* DNMTi – DNA methyltransferase inhibitor, HDACi – histone deacetylase inhibitor, ING (1–5) – inhibitor of growth (1–5), miR – microRNA, RSK2/CREB –serine/threonin kinase RSK2–cyclicAMP-response element-binding protein pathway, SAHA – suberoylanilide hydroxamic acid. *For details: see main text.*

showed synergistic effects with DNA-damaging agents and mitotic inhibitors to induce p53-mediated cell death in osteosarcoma cells [225,226]. Considering that *ING1* is downregulated in a variety of common human cancers, including breast cancer and GBM, and that it is a target of miR-622 in gastric cancer, studying the effects of modulating microRNA signatures that may regulate *ING1* in other malignancies may represent another step towards individualized cancer therapy.

*ING2* is the major molecular target of the HDACi suberoylanilide hydroxamic acid (SAHA) [227], suggesting that targeting of the INGs themselves may prove useful in combination with HDACi and other therapeutic agents that have been shown to induce *ING2*, thereby promoting p53-acetylation and subsequent apoptosis in cancer cells. This approach may also be effective in cancers upregulating *ING2*, such as gastrointestinal cancers. Intriguingly, *ING2* knockdown by siRNA enhanced growth suppression in adenocarcinoma cells most efficiently when combined with the widely used cytostatic adriamycin, whereas either single agent alone was significantly less effective [122]. As mentioned earlier, *ING2* knockdown suppresses MMP13, thus inhibiting tumor invasion [133,141]. Moreover, *ING2* knockdown sensitizes gastric cancer cells to chemosensitivity with 5-Fluoracil (5-FU) [141]. Therefore, therapy with siRNA against *ING2* may be an approach to patients with *ING2*-overexpressing tumors. However, methods of siRNA delivery to a targeted location still need to be optimized before going bedside. An interesting project would be to test, whether dose reduction of DNA-damaging agents, but adding SAHA to the drug regimen will result in similar or even higher efficacy. Also, inhibiting H3K4me3 binding by targeting the PHD, using small molecular compounds, may be a way to limit the oncogenic effects of *ING2*. The H3K4me3 binding region of the PHD forms substantial grooves [115], thereby representing potential targets for small molecules. However, this idea needs further refinement also, in particular with

regard to specificity towards the *ING2*, but not other PHD domains. Since not all *ING* proteins function as oncogenes, but do share the PHD, targeting this domain may not be broadly beneficial.

#### 2.10.2. Target-ING the MYST-ING3/4/5-HAT complexes

*ING3* is frequently downregulated in cancers, particularly in HNSCC and HCC. Moreover, *ING3* is negatively regulated by miR-21, a microRNA that is often upregulated in cancers and associated with tumor progression [118]. Hence, modulating miR-21 may increase *ING3*-expression and subsequently activate antiproliferative genes. In addition, RSK2 signaling through CREB downregulates *ING3*, thereby shielding HNSCC cells from apoptosis [160]. Thus, selective serine/threonin kinase inhibitors may increase *ING3* transcription and its transcriptionally activating effects.

In addition to various cancer hallmarks affected by all *ING* family proteins, *ING4* is the one that has been reported to additionally affect tumor promoting inflammation and immune destruction. Intriguingly, *ING4* has recently been suggested as a prognostic factor in breast cancer based on its ability to negatively regulate NF- $\kappa$ B target gene expression [184]. It also promoted apoptosis in lung cancer cells when co-expressed with Interleukin (IL)-20 [228]. *ING4* is silenced in breast, lung and many other human cancers, where it can promote sensitivity to conventional DNA damaging agents *in vitro* [229–232]. Considering that *ING4* is a target of miR-650 and miR-214 in some of these tumors, the possibility of combination therapy with tumor-specific miRNA modulators may be a promising approach.

This concept may also apply to cancers downregulating *ING5*, which is a target of miR-193. Interestingly, *ING5*-downregulation correlated with sensitivity to tamoxifen in ER-positive breast cancer cells in a genome-wide functional RNAi screen [233]. Thus, *ING5* might be involved in estrogen signaling in a subset of breast cancer patients. Moreover, *ING5* expression levels might serve as a

biomarker to recognize tamoxifen resistant breast cancer phenotypes. Considering, that the HBO1-JADE-HAT complex depends on its association with ING5 in order to enhance DNA replication, blocking ING5 by siRNA might be a future approach for the treatment of cancers overexpressing ING5, such as colorectal cancer.

#### Concluding Remarks

ING family proteins have proven to be potent targets for epigenetic cancer therapy, particularly in combination with conventional treatments. Their stoichiometric residence in HAT and HDAC complexes and links to DNA methylation make them logical choices for dysregulation of epigenetic status in cancer cells as a means of increasing the therapeutic index of many traditional cancer treatments.

#### Conflict of interest

None declared.

#### Acknowledgements

K.R. is a Scientist of the Alberta Heritage Foundation for Medical Research. This work was supported by the Canadian Institutes of Health Research and the Canadian Breast Cancer Foundation (to K.R.).

#### References

- [1] Hanahan, D. and Weinberg, R.A. (2011) Hallmarks of cancer: the next generation. *Cell* 144, 646–674.
- [2] Baylin, S.B. and Jones, P.A. (2011) A decade of exploring the cancer epigenome – biological and translational implications. *Nat. Rev. Cancer* 11, 726–734.
- [3] Vogelstein, B., Papadopoulos, N., Velculescu, V.E., et al. (2013) Cancer genome landscapes. *Science* 229, 1546–1558.
- [4] Berdasco, M. and Esteller, M. (2010) Aberrant epigenetic landscape in cancer: how cellular identity goes awry. *Dev. Cell* 19, 698–711.
- [5] Waddington, C.H. (2012) The epigenotype 1942. *Int. J. Epidemiol.* 41, 10–13.
- [6] Van Speybroeck, L. (2002) From epigenesis to epigenetics: the case of C.H. Waddington. *Ann. N.Y. Acad. Sci.* 981, 61–81.
- [7] Holliday, R. (2006) Epigenetics: a historical overview. *Epigenetics* 1, 76–80. Epub 2006 Mar 15.
- [8] Berger, S.L., Kouzarides, T., Shiekhattar, R., et al. (2009) An operational definition of epigenetics. *Genes Dev.* 23, 781–783.
- [9] Dawson, M.A. and Kouzarides, T. (2012) Cancer epigenetics: from mechanism to therapy. *Cell* 150, 12–27.
- [10] Jones, P.A. and Baylin, S.B. (2002) The fundamental role of epigenetic events in cancer. *Nat. Rev. Genet.* 3, 415–428.
- [11] Jones, P.A. and Baylin, S.B. (2007) The epigenomics of cancer. *Cell* 128, 683–692.
- [12] Bernstein, B.E., Mikkelsen, T.S., Xie, X., et al. (2006) A bivalent chromatin structure marks key developmental genes in embryonic stem cells. *Cell* 125, 315–326.
- [13] Bjerkvig, R., Tysnes, B.B., Aboody, K.S., et al. (2005) Opinion: the origin of the cancer stem cell: current controversies and new insights. *Nat. Rev. Cancer*, 899–904.
- [14] Bolden, J.E., Peart, M.J. and Johnstone, R.W. (2006) Anticancer activities of histone deacetylase inhibitors. *Nat. Rev. Drug Discovery* 5, 769–784.
- [15] Feinberg, A.P., Ohlsson, R. and Henikoff, S. (2006) The epigenetic progenitor origin of human cancer. *Nat. Rev. Genet.* 7, 21–33.
- [16] Guenther, M.G. and Young, R.A. (2010) Transcription. Repressive transcription. *Science* 329, 5988.
- [17] Brock, M.V., Hooker, C.M., Ota-Machida, E., et al. (2008) DNA methylation markers and early recurrence in stage I lung cancer. *N. Engl. J. Med.* 358, 1118–1128.
- [18] Mueller, W.C. and von Deimling, A. (2009) Gene regulation by methylation (von Deimling, Andreas, Ed.), *Gliomas, Recent Results in Cancer Research*, vol. 171, pp. 217–239, Springer, Berlin Heidelberg.
- [19] Saito, Y., Saito, H., Liang, G., et al. (2013) Epigenetic alterations and MicroRNA misexpression in cancer and autoimmune diseases: a critical review. *Clin. Rev. Allergy Immunol.* [Epub ahead of print].
- [20] Garkavtsev, I., Kazarov, A., Gudkov, A., et al. (1996) Suppression of the novel growth inhibitor p33ING1 promotes neoplastic transformation. *Nat. Genet.* 14, 415–420.
- [21] He, G.H., Helbing, C.C., Wagner, M.J., et al. (2005) Phylogenetic analysis of the ING family of PHD finger proteins. *Mol. Biol. Evol.* 22, 104–116.
- [22] Boveri, T. (1914) Zur Frage der Entstehung Maligner Tumoren, G. Fischer.
- [23] Reddy, K.L. and Feinberg, A.P. (2013) Higher order chromatin organization in cancer. *Semin. Cancer Biol.* 23, 109–115.
- [24] Satpathy, S., Nabbi, A. and Riabowol, K. (2013) Regulation of chromatin regulators: post-translational modification of the ING family of epigenetic regulators. *Biochem. J.* 450, 433–442.
- [25] Shi, X., Hong, T., Walter, K.L., et al. (2006) ING2 PHD domain links histone H3 lysine 4 methylation to active gene repression. *Nature* 442, 96–99.
- [26] Li, H., Ilin, S., Wang, W., et al. (2006) Molecular basis for site-specific read-out of histone H3K4me3 by the BPTF PHD finger of NURF. *Nature* 442, 91–95.
- [27] Liu, Y., Subrahmanyam, R., Chakraborty, T., et al. (2007) A plant homeodomain in RAG-2 that binds hypermethylated lysine 4 of histone H3 is necessary for efficient antigen-receptor-gene rearrangement. *Immunity* 27, 561–571.
- [28] Matthews, A.G., Kuo, A.J., Ramon-Maiques, S., et al. (2007) RAG2 PHD finger couples histone H3 lysine 4 trimethylation with V(D)J recombination. *Nature* 450, 1106–1110.
- [29] Pefia, P.V., Davrazou, F., Shi, X., et al. (2006) Molecular mechanism of histone H3K4me3 recognition by plant homeodomain of ING2. *Nature* 442, 100–103.
- [30] Ramon-Maiques, S., Kuo, A.J., Carney, D., et al. (2007) The plant homeodomain finger of RAG2 recognizes histone H3 methylated at both lysine-4 and arginine-2. *Proc. Natl. Acad. Sci. U.S.A.* 104, 18993–18998.
- [31] Ruthenburg, A.J., Allis, C.D. and Wysocka, J. (2007) Methylation of lysine 4 on histone H3: intricacy of writing and reading a single epigenetic mark. *Mol. Cell* 25, 15–30.
- [32] Saksouk, N., Awakumov, N. and Cote, J. (2008) (de)MYSTification and INGenuity of tumor suppressors. *Cell. Mol. Life Sci.* 65, 1013–1018.
- [33] Vermeulen, M., Mulder, K.W. and Denissov, S. et al. (2007) Selective anchoring of TFIID to nucleosomes by trimethylation of histone H3 lysine 4. *Cell* 131, 58–69.
- [34] Wysocka, J., Swigut, T., Xiao, H., et al. (2006) A PHD finger of NURF couples histone H3 lysine 4 trimethylation with chromatin remodelling. *Nature* 442, 86–90.
- [35] Taverna, S.D., Ilin, S., Rogers, R.S., et al. (2006) Yng1 PHD finger binding to H3 trimethylated at K4 promotes NuA3 HAT activity at K14 of H3 and transcription at a subset of targeted ORFs. *Mol. Cell* 24, 785–796.
- [36] Hung, T., Binda, O., Champagne, S., et al. (2009) ING4 mediates crosstalk between histone H3 K4 trimethylation and H3 acetylation to attenuate cellular transformation. *Mol. Cell* 33, 248–256.
- [37] Saksouk, N., Awakumov, N., Champagne, K.S., et al. (2009) HBO1 HAT complexes target chromatin throughout gene coding regions via multiple PHD finger interactions with histone H3 tail. *Mol. Cell* 33, 257–265.
- [38] Schafer, A., Karaulanov, E., Stapf, U., et al. (2013) Ing1 functions in DNA demethylation by directing Gadd45a to H3K4me3. *Genes Dev.* 27, 261–273.
- [39] Schomacher, L. (2013) Mammalian DNA demethylation: multiple faces and upstream regulation. *Epigenetics* 8, 679–684.
- [40] Russell, M., Berardi, P., Gong, W., et al. (2006) Grow-ING, Age-ING and Die-ING: ING proteins link cancer, senescence and apoptosis. *Exp. Cell Res.* 312, 951–961.
- [41] Scott, M., Boisvert, F.M., Vieyra, D., et al. (2001) UV induces nucleolar translocation of ING1 through two distinct nucleolar targeting sequences. *Nucleic Acids Res.* 29, 2052–2058.
- [42] Eriksson, M., Brown, W.T., Gordon, L.B., et al. (2003) Recurrent *de novo* point mutations in lamin A cause Hutchinson–Gilford progeria syndrome. *Nature* 423, 293–298.
- [43] Han, X., Feng, X., Rattner, J.B., et al. (2008) Tethering by lamin A stabilizes and targets the ING1 tumour suppressor. *Nat. Cell Biol.* 10, 1333–1340.
- [44] Gilbert, S.F. (2009) Ageing and cancer as diseases of epigenesis. *J. Biosci.* 34, 601–604.
- [45] Espada, J., Ballestar, E., Santoro, R., et al. (2007) Epigenetic disruption of ribosomal RNA genes and nucleolar architecture in DNA methyltransferase 1 (Dnmt1) deficient cells. *Nucleic Acids Res.* 35, 2191–2198.
- [46] Agrelo, R., Setien, F., Espada, J., et al. (2005) Inactivation of the lamin A/C gene by CpG island promoter hypermethylation in hematologic malignancies: and its association with poor survival in nodal diffuse large B-cell lymphoma. *J. Clin. Oncol.* 23, 3940–3947.
- [47] Bjornsson, H.T., Sigurdsson, M.I., Fallin, M.D., et al. (2008) Intra-individual change over time in DNA methylation with familial clustering. *JAMA J. Am. Medical Assoc.* 299, 2877–2883.
- [48] Stadelmann, B., Khandjian, E., Hirt, A., et al. (1990) Repression of nuclear lamin A and C gene expression in human acute lymphoblastic leukemia and non-Hodgkin's lymphoma cells. *Leukemia Res.* 4, 815–821.
- [49] Nouman, G.S., Anderson, J.J., Crosier, S., et al. (2002) Loss of nuclear expression of the p33(ING1b) inhibitor of growth protein in childhood acute lymphoblastic leukaemia. *J. Clin. Pathol.* 55, 596–601.
- [50] Kumamoto, K., Fujita, R., Kuratani, M., et al. (2009) ING2 is upregulated in colon cancer and increases invasion by enhanced MMP13 expression. *Int. J. Cancer* 125, 1306–1315.
- [51] Belt, E.J.T., Fijneman, R.J., van den Berg, E.G., et al. (2011) Loss of lamin A/C expression in stage II and III colon cancer is associated with disease recurrence. *Eur. J. Cancer* 47, 1837–1845.
- [52] Kuzmichev, A., Zhang, Y., Erdjument-Bromage, H., et al. (2002) Role of the Sin3–histone deacetylase complex in growth regulation by the candidate tumor suppressor p33(ING1). *Mol. Cell Biol.* 22, 835–848.



- [53] Doyon, Y., Cayrou, C., Ullah, M., et al. (2006) ING tumor suppressor proteins are critical regulators of chromatin acetylation required for genome expression and perpetuation. *Mol. Cell* 21, 51–64.
- [54] Gozani, O., Karuman, P., Jones, D.R., et al. (2003) The PHD finger of the chromatin-associated protein ING2 functions as a nuclear phosphoinositide receptor. *Cell* 114, 99–111.
- [55] Kaadige, M.R. and Ayer, D.E. (2006) The polybasic region that follows the plant homeodomain zinc finger 1 of Pfl is necessary and sufficient for specific phosphoinositide binding. *J. Biol. Chem.* 281, 28831–28836.
- [56] Thalappilly, S., Feng, X., Pastyrkova, S., et al. (2011) The p53 tumor suppressor is stabilized by inhibitor of growth 1 (ING1) by blocking polyubiquitination. *PLoS One* 6, 28831–28836.
- [57] Eapen, S.A., Netherton, S.J., Sarker, K.P., et al. (2012) Identification of a novel function for the chromatin remodeling protein ING2 in muscle differentiation. *PLoS One* 7, e40684.
- [58] Scott, M., Bonnefin, P., Vieyra, D., et al. (2001) UV-induced binding of ING1 to PCNA regulates the induction of apoptosis. *J. Cell Sci.* 114, 3455–3462.
- [59] Wong, R.P., Lin, H., Khosravi, S., et al. (2011) Tumour suppressor ING1b maintains genomic stability upon replication stress. *Nucleic Acids Res.* 39, 3632–3642.
- [60] Satpathy, S., Guerillon, C., Thakur, S., et al., SUMOylation of the ING1 tumour suppressor affects transcriptional repression. *Carcinogenesis*, in press.
- [61] Unoki, M., Kumamoto, K. and Harris, C.C. (2009) ING proteins as anticancer drugs. *Curr. Drug Targets* 10, 442–454.
- [62] Feng, X., Hara, Y. and Riabowol, K. (2002) Different HATS of the ING1 gene family. *Trends Cell Biol.* 12, 532–538.
- [63] Soliman, M.A. and Riabowol, K. (2007) After a decade of study-ING, a PHD for a versatile family of proteins. *Trends Biochem. Sci.* 32, 509–519.
- [64] Soliman, M.A., Berardi, P., Pastyrkova, S., et al. (2008) ING1a expression increases during replicative senescence and induces a senescent phenotype. *Aging Cell* 7, 783–794.
- [65] Rajarajacholan, U.K., Thalappilly, S. and Riabowol, K. (2013) The ING1a tumor suppressor regulates endocytosis to induce cellular senescence via the Rb-E2F pathway. *PLoS Biol.* 11, e1001502.
- [66] Coles, A.H., Liang, H., Zhu, Z., et al. (2007) Deletion of p37Ing1 in mice reveals a p53-independent role for Ing1 in the suppression of cell proliferation, apoptosis, and tumorigenesis. *Cancer Res.* 67, 2054–2061.
- [67] Luo, J., Shah, S., Riabowol, K., et al. (2009) The *Caenorhabditis elegans* ing-3 gene regulates ionizing radiation-induced germ-cell apoptosis in a p53-associated pathway. *Genetics* 181, 473–482.
- [68] Helbing, C.C., Veillette, C., Riabowol, K., et al. (1997) A novel candidate tumor suppressor, ING1, is involved in the regulation of apoptosis. *Cancer Res.* 57, 1255–1258.
- [69] Bose, P., Thakur, S. and Thalappilly, S. (2013) ING1 induces apoptosis through direct effects at the mitochondria. *Cell Death Dis.* 4, e788.
- [70] Takahashi, M., Ozaki, T., Todo, S., et al. (2004) Decreased expression of the candidate tumor suppressor gene ING1 is associated with poor prognosis in advanced neuroblastomas. *Oncol. Rep.* 12, 811–816.
- [71] Campos, E.I., Cheung Jr., K.J., Murray, A., et al. (2002) The novel tumour suppressor gene ING1 is overexpressed in human melanoma cell lines. *Br. J. Dermatol.* 146, 574–580.
- [72] Stark, M., Puig-Butille, J.A., Walker, G., et al. (2006) Mutation of the tumour suppressor p33ING1b is rare in melanoma. *Br. J. Dermatol.* 155, 94–99.
- [73] Nouman, G.S., Anderson, J.J., Mathers, M.E., et al. (2002) Nuclear to cytoplasmic shift of the p33(ING1b) tumour suppressor protein is associated with malignancy in melanocytic lesions. *Histopathology* 40, 360–366.
- [74] Campos, E.I., Martinka, M., Mitchell, D.L., et al. (2004) Mutations of the ING1 tumor suppressor gene detected in human melanoma abrogate nucleotide excision repair. *Int. J. Oncol.* 25, 73–80.
- [75] Walzak, A.A., Veldhoen, N., Feng, X., et al. (2008) Expression profiles of mRNA transcript variants encoding the human inhibitor of growth tumor suppressor gene family in normal and neoplastic tissues. *Exp. Cell Res.* 314, 273–285.
- [76] Kameyama, K., Huang, C.L., Liu, D., et al. (2003) Reduced ING1b expression plays an important role in carcinogenesis of non-small cell lung cancer patients. *Clin. Cancer Res.* 9, 5952–5961. Liu, J., Lin, Y., Yang, H., et al. (2011) The expression of p33(ING1), p53, and autophagy-related gene Beclin1 in patients with non-small cell lung cancer. *Tumour Biol.* 32, 1113–1121.
- [77] Shen, D.H., Chan, K.Y., Khoo, U.S., et al. (2005) Epigenetic and genetic alterations of p33ING1b in ovarian cancer. *Carcinogenesis* 26, 855–863.
- [78] Liu, J., Lin, Y., Yang, H., et al. (2011) The expression of p33(ING1), p53, and autophagy-related gene Beclin1 in patients with non-small cell lung cancer. *Tumour Biol.* 32, 1113–1121.
- [79] Vieyra, D., Senger, D.L., Toyama, T., et al. (2003) Altered subcellular localization and low frequency of mutations of ING1 in human brain tumors. *Clin. Cancer Res.* 9, 5952–5961.
- [80] Tallen, G., Kaiser, I., Krabbe, S., et al. (2004) No ING1 mutations in human brain tumours but reduced expression in high malignancy grades of astrocytoma. *Int. J. Cancer* 109, 476–479.
- [81] Oki, E., Maehara, Y., Tokunaga, E., et al. (1999) Reduced expression of p33(ING1) and the relationship with p53 expression in human gastric cancer. *Cancer Lett.* 147, 157–162.
- [82] Chen, L.S., Wei, J.B., Zhou, Y.C., et al. (2005) Genetic alterations and expression of inhibitor of growth 1 in human sporadic colorectal cancer. *World J. Gastroenterol.* 11, 6120–6124.
- [83] Chen, L., Matsubara, N., Yoshino, T., et al. (2001) Genetic alterations of candidate tumor suppressor ING1 in human esophageal squamous cell cancer. *Cancer Res.* 61, 4345–4349.
- [84] Zhu, Z., Lin, J., Qu, J.H., et al. (2005) Inhibitory effect of tumor suppressor p33(ING1b) and its synergy with p53 gene in hepatocellular carcinoma. *World J. Gastroenterol.* 11, 1903–1909.
- [85] Yu, G.Z., Zhu, M.H., Zhu, Z., et al. (2004) Genetic alterations and reduced expression of tumor suppressor p33(ING1b) in human exocrine pancreatic carcinoma. *World J. Gastroenterol.* 10, 3597–3601.
- [86] Tokunaga, E., Maehara, Y., Oki, E., et al. (2000) Diminished expression of ING1 mRNA and the correlation with p53 expression in breast cancers. *Cancer Lett.* 2000 152, 15–22.
- [87] Nouman, G.S., Anderson, J.J., Crosier, S., et al. (2003) Downregulation of nuclear expression of the p33(ING1b) inhibitor of growth protein in invasive carcinoma of the breast. *J. Clin. Pathol.* 56, 507–511.
- [88] Jager, D., Stockert, E., Scanlan, M.J., et al. (1999) Cancer-testis antigens and ING1 tumor suppressor gene product are breast cancer antigens: characterization of tissue-specific ING1 transcripts and a homologue gene. *Cancer Res.* 59, 6197–6204.
- [89] Thakur, S., Feng, X., Qiao Shi, Z., et al. (2012) ING1 and 5-azacytidine act synergistically to block breast cancer cell growth. *PLoS One* 7, e43671.
- [90] Sanchez-Carbayo, M., Socci, N.D., Lozano, J.J., et al. (2003) Gene discovery in bladder cancer progression using cDNA microarrays. *Am. J. Pathol.* 163, 505–516.
- [91] Mihara, M., Erster, S., Zaika, A., et al. (2003) p53 Has a direct apoptogenic role at the mitochondria. *Mol. Cell* 11 (3), 577–590. 2003 Mar.
- [92] Kichina, J.V., Zeremski, M., Aris, L., et al. (2006) Targeted disruption of the mouse *ingl* locus results in reduced body size, hypersensitivity to radiation and elevated incidence of lymphomas. *Oncogene* 25, 857–866.
- [93] Saito, M., Kumamoto, K., Robles, A.I., et al. (2010) Targeted disruption of *Ing2* results in defective spermatogenesis and development of soft-tissue sarcomas. *PLoS One* 5, e15541.
- [94] Coles, A.H., Gannon, H., Cerny, A., et al. (2010) Inhibitor of growth-4 promotes IκB promoter activation to suppress NF-κB signaling and innate immunity. *Proc. Natl. Acad. Sci. U.S.A.* 107, 11423–11428.
- [95] Shah, S. and Riabowol, K. (2009) Signaling pathways of the ING proteins in apoptosis. *Curr. Drug Targets* 10, 385–391.
- [96] Gunduz, M., Ouchida, M., Fukushima, K., et al. (2000) Genomic structure of the human ING1 gene and tumor-specific mutations detected in head and neck squamous cell carcinomas. *Cancer Res.* 60, 3143–3146.
- [97] Jones, P.A. (2012) Functions of DNA methylation: islands, start sites, gene bodies and beyond. *Nat. Rev. Genet.* 13, 484–492.
- [98] Tong, W.G., Wierda, W.G., Lin, E., et al. (2010) Genome-wide DNA methylation profiling of chronic lymphocytic leukemia allows identification of epigenetically repressed molecular pathways with clinical impact. *Epigenetics* 5, 499–508.
- [99] Zhu, X. and Zeisel, S.H. (2005) Gene expression profiling in phosphatidylethanolamine N-methyltransferase knockout mice. *Brain Res. Mol. Brain Res.* 134, 239–255.
- [100] Guo, X.B., Jing, C.Q., Li, L.P., et al. (2011) Down-regulation of miR-622 in gastric cancer promotes cellular invasion and tumor metastasis by targeting ING1 gene. *World J. Gastroenterol.* 17, 1895–1902.
- [101] Garate, M., Campos, E.I., Bush, J.A., et al. (2007) Phosphorylation of the tumor suppressor p33(ING1b) at Ser-126 influences its protein stability and proliferation of melanoma cells. *FASEB J.* 21, 3705–3716.
- [102] Garate, M., Wong, R.P., Campos, E.I., et al. (2008) NAD(P)H quinone oxidoreductase 1 inhibits the proteasomal degradation of the tumour suppressor p33(ING1b). *EMBO Rep.* 9, 576–581.
- [103] Manning, G., Whyte, D.B., Martinez, R., et al. (2002) The protein kinase complement of the human genome. *Science* 298, 1912–1934.
- [104] Yu, L., Thakur, S., Leong-Quong, R.Y., et al. (2013) Src regulates the activity of the ING1 tumor suppressor. *PLoS One* 8, e60943.
- [105] Abad, M., Menendez, C., Fuchtbauer, A., et al. (2007) Ing1 mediates p53 accumulation and chromatin modification in response to oncogenic stress. *J. Biol. Chem.* 282, 31060–31067.
- [106] Helbing, C.C., Wagner, M.J., Pettem, K., et al. (2011) Modulation of thyroid hormone-dependent gene expression in *Xenopus laevis* by inhibitor of growth (ING) proteins. *PLoS One* 6, e28658.
- [107] Tallen, G., Farhangi, S., Tamannai, M., et al. (2011) The inhibitor of growth 1 (ING1) proteins suppress angiogenesis and differentially regulate angiopoietin expression in glioblastoma cells. *PLoS One* 6, e21065.
- [108] Vieyra, D., Toyama, T., Hara, Y., et al. (2002) ING1 isoforms differentially affect apoptosis in a cell age-dependent manner. *Cancer Res.* 62, 4445–4452.
- [109] Shinoura, N., Muramatsu, Y., Nishimura, M., et al. (1999) Adenovirus-mediated transfer of p33ING1 with p53 drastically augments apoptosis in gliomas. *Cancer Res.* 59, 5521–5528.
- [110] Coles, A.H., Marfella, C.G., Imbalzano, A.N., et al. (2008) p37Ing1b regulates B-cell proliferation and cooperates with p53 to suppress diffuse large B-cell lymphomagenesis. *Cancer Res.* 68, 8705–8714.
- [111] Ceruti, J.M., Ogara, M.F., Menendez, C., et al. (2013) Inhibitor of growth 1 (ING1) acts at early steps of multiple DNA repair pathways. *Mol. Cell. Biochem.* 378, 117–126.
- [112] Skowyrda, D., Zeremski, M., Neznanov, N., et al. (2012) Differential association of products of alternative transcripts of the candidate tumor suppressor ING1 with the mSin3/HDAC1 transcriptional corepressor complex. *J. Biol. Chem.* 276, 8734–8739.



- [113] Goeman, F., Thormeyer, D., Abad, M., et al. (2005) Growth inhibition by the tumor suppressor p33ING1 in immortalized and primary cells: involvement of two silencing domains and effect of Ras. *Mol. Cell. Biol.* 25, 422–431.
- [114] Vieyra, D., Loewith, R., Scott, M., et al. (2002) Human ING1 proteins differentially regulate histone acetylation. *J. Biol. Chem.* 277, 832–839.
- [115] Li, N., Li, Q., Cao, X., et al. (2011) The tumor suppressor p33ING1b upregulates p16INK4a expression and induces cellular senescence. *FEBS Lett.* 585, 3106–3112.
- [116] Xin, H., Yoon, H.G., Singh, P.B., et al. (2004) Components of a pathway maintaining histone modification and heterochromatin protein 1 binding at the pericentric heterochromatin in mammalian cells. *J. Biol. Chem.* 279, 9539–9546.
- [117] Chen, J., Tran, U.M., Rajarajacholan, U., et al. (2013) ING1b-inducible microRNA203 inhibits cell proliferation. *Br. J. Cancer* 108, 1143–1148.
- [118] Gomez-Cabello, D., Callejas, S., Benguria, A., et al. (2010) Regulation of the microRNA processor DGCR8 by the tumor suppressor ING1. *Cancer Res.* 70, 1866–1874.
- [119] Shimada, Y., Saito, A., Suzuki, M., et al. (1998) Cloning of a novel gene (ING1L) homologous to ING1, a candidate tumor suppressor. *Cytogenet. Cell Genet.* 83, 232–235.
- [120] Nagashima, M., Shiseki, M., Miura, K., et al. (2001) DNA damage- inducible gene p33ING2 negatively regulates cell proliferation through acetylation of p53. *Proc. Natl. Acad. Sci. U.S.A.* 98, 9671–9676.
- [121] Unoki, M., Kumamoto, K., Robles, A.I., et al. (2008) A novel ING2 isoform, ING2b, synergizes with ING2a to prevent cell cycle arrest and apoptosis. *FEBS Lett.* 582, 3868–3874.
- [122] Wang, Y., Wang, J. and Li, G. (2006) Leucine zipper-like domain is required for tumor suppressor ING2-mediated nucleotide excision repair and apoptosis. *FEBS Lett.* 580, 3787–3793.
- [123] Gozani, O., Karuman, P., Jones, D.R., et al. (2003) The PHD finger of the chromatin-associated protein ING2 functions as a nuclear phosphoinositide receptor. *Cell* 114, 99–111.
- [124] Eapen, S.A., Netherton, S.J., Sarker, K.P., et al. (2012) Identification of a novel function for the chromatin remodeling protein ING2 in muscle differentiation. *PLoS One* 7, e40684.
- [125] Sarker, K.P., Kataoka, H., Chan, A., et al. (2008) ING2 as a novel mediator of transforming growth factor-beta-dependent responses in epithelial cells. *J. Biol. Chem.* 283, 13269–13279.
- [126] Zhang, H.K., Pan, K., Wang, H., et al. (2008) Decreased expression of ING2 gene and its clinicopathological significance in hepatocellular carcinoma. *Cancer Lett.* 261, 183–192.
- [127] Zhang, H., Ma, H., Wang, Q., et al. (2010) Analysis of loss of heterozygosity on chromosome 4q in hepatocellular carcinoma using high-throughput SNP array. *Oncol. Rep.* 23, 445–455.
- [128] Borkosky, S.S., Gunduz, M., Nagatsuka, H., et al. (2009) Frequent deletion of ING2 locus at 4q35.1 associates with advanced tumor stage in head and neck squamous cell carcinoma. *J. Cancer Res. Clin. Oncol.* 135, 703–713.
- [129] Okano, T., Gemma, A., Hosoya, Y., et al. (2006) Alterations in novel candidate tumor suppressor genes, ING1 and ING2 in human lung cancer. *Oncol. Rep.* 15, 545–549.
- [130] Ythier, D., Brambilla, E., Binet, R., et al. (2010) Expression of candidate tumor suppressor gene ING2 is lost in non-small cell lung carcinoma. *Lung Cancer* 69, 180–186.
- [131] Kumamoto, K., Spillare, E.A., Fujita, K., et al. (2008) Nutlin-3a activates p53 to both down-regulate inhibitor of growth 2 and up-regulate mir-34a, mir-34b, and mir-34c expression, and induce senescence. *Cancer Res.* 68, 3193–3203.
- [132] Meulmeester, E. and Jochemsen, A.G. (2008) p53: a guide to apoptosis. *Curr. Cancer Drug Targets* 8, 87–97.
- [133] Kumamoto, K., Fujita, K. and Kurotani, R. (2009) ING2 is upregulated in colon cancer and increases invasion by enhanced MMP13 expression. *Int. J. Cancer* 125, 1306–1315.
- [134] Nie, J., Liu, L., Wu, M., et al. (2010) HECT ubiquitin ligase Smurf1 targets the tumor suppressor ING2 for ubiquitination and degradation. *FEBS Lett.* 584, 3005–3012.
- [135] Ythier, D., Larrieu, D., Binet, R., et al. (2010) Sumoylation of ING2 regulates the transcription mediated by Sin3A. *Oncogene* 29, 5946–5956.
- [136] Pedoux, R., Sengupta, S., Shen, J.C., et al. (2005) ING2 regulates the onset of replicative senescence by induction of p300-dependent p53 acetylation. *Mol. Cell. Biol.* 25, 6639–6648.
- [137] Larrieu, D., Ythier, D., Brabilla, C., et al. (2010) ING2 controls the G1 to S-phase transition by regulating p21 expression. *Cell Cycle* 9, 3984–3990.
- [138] Sarker, K.P., Kataoka, H., Chan, A., et al. (2008) ING2 as a novel mediator of transforming growth factor-beta-dependent responses in epithelial cells. *J. Biol. Chem.* 283, 269–279.
- [139] Roberts, A.B. and Wakefield, L.M. (2003) The two faces of transforming growth factor beta in carcinogenesis. *Proc. Natl. Acad. Sci. U.S.A.* 100, 8621–8623.
- [140] Wang, J., Chin, M.Y. and Li, G. (2006) The novel tumor suppressor p33ING2 enhances nucleotide excision repair via induction of histone H4 acetylation and chromatin relaxation. *Cancer Res.* 66, 1906–1911.
- [141] Zhong, J., Yang, L., Liu, N., et al. (2013) Knockdown of inhibitor of growth protein 2 inhibits cell invasion and enhances chemosensitivity to 5-FU in human gastric cancer cells. *Dig. Dis. Sci.* 58, 3189–3197.
- [142] Pena, P.V., Davrazou, F., Shi, X., et al. (2006) Molecular mechanism of histone H3K4me3 recognition by plant homeodomain of ING2. *Nature* 442, 100–103.
- [143] Shi, X., Hong, T., Walter, K.L., et al. (2006) ING2 PHD domain links histone H3 lysine 4 methylation to active gene repression. *Nature* 442, 96–99.
- [144] Bua, D.J., Martin, G.M., Binda, O., et al. (2013) Nuclear phosphatidylinositol-5-phosphate regulates ING2 stability at discrete chromatin targets in response to DNA damage. *Sci. Rep.* 3, 2137.
- [145] Leeman, M.F., Curran, S. and Murray, G.I. (2002) The structure, regulation, and function of human matrix metalloproteinase-13.143. *Crit. Rev. Biochem. Mol. Biol.* 37 (3), 149–166.
- [146] Lai, A., Kennedy, B.K., Barbie, D.A., et al. (2001) RBP1 recruits the mSIN3-histone deacetylase complex to the pocket of retinoblastoma tumor suppressor family proteins found in limited discrete regions of the nucleus at growth arrest. *Mol. Cell. Biol.* 21, 2918–2932.
- [147] Binda, O., Nassif, C. and Branton, P.E. (2008) SIRT1 negatively regulates HDAC1-dependent transcriptional repression by the RBP1 family of proteins. *Oncogene* 27, 3384–3392.
- [148] Fegers, I., Kob, R., Eckey, M., et al. (2007) The tumor suppressors p33ING1 and p33ING2 interact with alien in vivo and enhance alien-mediated gene silencing. *J. Proteome Res.* 6, 4182–4188.
- [149] Goeman, F., Otto, K., Kyrilenko, S., Schmidt, O. and Baniahmad, A. (2008) ING2 recruits histone methyltransferase activity with methylation site specificity distinct from histone H3 lysines 4 and 9. *Biochim. Biophys. Acta* 1783, 1673–1680.
- [150] Nagashima, M., Shiseki, M., Pedoux, R.M., et al. (2003) A novel PHD- finger motif protein, p47ING3, modulates p53-mediated transcription, cell cycle control, and apoptosis. *Oncogene* 22, 343–350.
- [151] Wang, Y. and Li, G. (2006) ING3 promotes UV-induced apoptosis via Fas/ caspase-8 pathway in melanoma cells. *J. Biol. Chem.* 281, 887–893.
- [152] Awe, J.P. and Byrne, J.A. (2013) Identifying candidate oocyte reprogramming factors using cross-species global transcriptional analysis. *Cell Reprogram.* 15, 126–133.
- [153] Luo, J., Shah, S., Riabowol, K., et al. (2009) The *Caenorhabditis elegans* ing-3 gene regulates ionizing radiation-induced germ-cell apoptosis in a p53-associated pathway. *Genetics* 181, 473–482.
- [154] Suzuki, S., Nozawa, Y., Tsukamoto, S., et al. (2013) ING3 is essential for asymmetric cell division during mouse oocyte maturation. *PLoS One* 8, e74749.
- [155] Cekaite, L., Clancy, T. and Sioud, M. (2010) Increased miR-21 expression during human monocyte differentiation into DCs. *Front Biosci. (Elite Ed)* 2, 818–828.
- [156] Gunduz, M., Ouchida, M., Fukushima, K., et al. (2002) Allelic loss and reduced expression of the ING3, a candidate tumor suppressor gene at 7q31, in human head and neck cancers. *Oncogene* 21, 4462–4470.
- [157] Gunduz, M., Beder, L.B., Gunduz, E., et al. (2008) Downregulation of ING3 mRNA expression predicts poor prognosis in head and neck cancer. *Cancer Sci.* 99, 531–538.
- [158] Lu, M., Chen, F., Wang, Q., et al. (2012) Downregulation of inhibitor of growth 3 is correlated with tumorigenesis and progression of hepatocellular carcinoma. *Oncol. Lett.* 4, 47–52.
- [159] Yang, H.Y., Liu, H.L., Tian, L.T., et al. (2012) Expression and prognostic value of ING3 in human primary hepatocellular carcinoma. *Exp. Biol. Med.* (Maywood) (237), 352–361.
- [160] Jin, L., Li, D., Lee, J.S., et al. (2013) p90 RSK2 mediates antianoinis signals by both transcription-dependent and -independent mechanisms. *Mol. Cell. Biol.* 33, 2574–2585.
- [161] Choudhary, C., Kumar, C., Gnäd, F., et al. (2009) Lysine acetylation targets protein complexes and co-regulates major cellular functions. *Science* 325, 834–840.
- [162] Chen, G., Wang, Y., Garate, M., et al. (2010) The tumor suppressor ING3 is degraded by SCF(Skp2)-mediated ubiquitin-proteasome system. *Oncogene* 29, 1498–1508.
- [163] Doyon, Y., Selleck, W., Lane, W.S., et al. (2004) Structural and functional conservation of the NuA4 histone acetyltransferase complex from yeast to humans. *Mol. Cell. Biol.* 24, 1884–1896.
- [164] Awakumov, N. and Cote, J. (2007) 162 The MYST family of histone acetyltransferases and their intimate links to cancer. *Oncogene* 26, 5395–5407.
- [165] Squatrito, M., Gorriani, C. and Amati, B. (2006) Tip60 in DNA damage response and growth control: many tricks in one HAT. *Trends Cell Biol.* 16, 433–442.
- [166] Unoki, M., Shen, J.C., Zheng, Z.M., et al. (2006) Novel splice variants of ING4 and their possible roles in the regulation of cell growth and motility. *J. Biol. Chem.* 281, 677–686.
- [167] Raho, G., Miranda, C., Tamborini, E., et al. (2007) Detection of novel mRNA splice variants of human ING4 tumor suppressor gene. *Oncogene* 26, 5247–5257.
- [168] Kim, S., Chin, K., Gray, J.W., et al. (2004) A screen for genes that suppress loss of contact inhibition: identification of ING4 as a candidate tumor suppressor gene in human cancer. *Proc. Natl. Acad. Sci. U.S.A.* 101, 16251–16256.
- [169] Zhang, X., Wang, K.S., Wang, Z.Q., et al. (2005) Nuclear localization signal of ING4 plays a key role in its binding to p53. *Biochem. Biophys. Res. Commun.* 331, 1032–1038.
- [170] Tsai, K.W., Tseng, H.C. and Lin, W.C. (2008) Two wobble-splicing events affect ING4 protein subnuclear localization and degradation. *Exp. Cell Res.* 314, 3130–3141.
- [171] Li, X., Cai, L., Chen, H., et al. (2009) Inhibitor of growth 4 induces growth suppression and apoptosis in glioma U87MG. *Pathobiology* 76, 181–192.
- [172] Kiriomonos, G., Bravou, V., Papachristou, D.J., et al. (2010) Loss of inhibitor of growth (ING-4) is implicated in the pathogenesis and progression of human astrocytomas. *Brain Pathol.* 20, 490–497.

- [173] Kim, S., Welm, A.L. and Bishop, J.M. (2010) A dominant mutant allele of the ING4 tumor suppressor found in human cancer cells exacerbates MYC-initiated mouse mammary tumorigenesis. *Cancer Res.* 70, 5155–5162.
- [174] Byron, S.A., Min, E., Thai, T.S., et al. (2012) Negative regulation of NF- $\kappa$ B by the ING4 tumor suppressor in breast cancer. *PLoS One* 7, e46823.
- [175] Zhang, L., Wang, Y., Zhang, F., et al. (2012) Correlation between tumor suppressor inhibitor of growth family member 4 expression and microvessel density in breast cancer. *Hum. Pathol.* 43, 1611–1617.
- [176] Li, M., Jin, Y., Sun, W.J., et al. (2009) Reduced expression and novel splice variants of ING4 in human gastric adenocarcinoma. *J. Pathol.* 219, 87–95.
- [177] Nanding, A., Tang, L., Cai, L., et al. (2014) Low ING4 protein expression detected by paraffin-section immunohistochemistry is associated with poor prognosis in untreated patients with gastrointestinal stromal tumors. *Gastric Cancer* 17, 87–96.
- [178] Li, S., Fan, T., Liu, H., et al. (2013) Tumor suppressor ING4 overexpression contributes to proliferation and invasion inhibition in gastric carcinoma by suppressing the NF- $\kappa$ B signaling pathway. *Mol. Biol. Rep.* 40, 5723–5732.
- [179] Gunduz, M., Nagatsuka, H., Demircan, K., et al. (2005) Frequent deletion and down-regulation of ING4, a candidate tumor suppressor gene at 12p13, in head and neck squamous cell carcinomas. *Gene* 356, 109–117.
- [180] Li, X.H., Kikuchi, K., Zheng, Y., et al. (2011) Downregulation and translocation of nuclear ING4 is correlated with tumorigenesis and progression of head and neck squamous cell carcinoma. *Oral Oncol.* 47, 217–223.
- [181] Zeng, Z.L., Li, F.J., Gao, F., et al. (2013) Upregulation of miR-650 is correlated with down-regulation of ING4 and progression of hepatocellular carcinoma. *J. Surgical Oncol.* 107, 105–110.
- [182] Fang, F., Luo, L.B., Tao, Y.M., et al. (2009) Decreased expression of inhibitor of growth 4 correlated with poor prognosis of hepatocellular carcinoma. *Cancer Epidemiol. Biomarkers Prev.* 18, 409–416.
- [183] Cai, L., Li, X., Zheng, S., et al. (2009) Inhibitor of growth 4 is involved in melanomagenesis and induces growth suppression and apoptosis in melanoma cell line M14. *Melanoma Res.* 19, 1–7.
- [184] Wei, Q., He, W., Lu, Y., et al. (2012) Effect of the tumor suppressor gene ING4 on the proliferation of MCF-7 human breast cancer cells. *Oncol. Lett.* 4, 438–442.
- [185] Lu, J., Tang, Y., Chen, Y., et al. (2013) ING4 regulates JWA in angiogenesis and their prognostic value in melanoma patients. *Br. J. Cancer* 109, 2842–2852.
- [186] Lou, C., Jiang, S., Guo, X., et al. (2012) ING4 is negatively correlated with microvessel density in colon cancer. *Tumour Biol.* 33, 2357–2364.
- [187] Liu, Y., Yu, L., Wang, Y., et al. (2012) Expression of tumor suppressor gene ING4 in ovarian carcinoma is correlated with microvessel density. *J. Cancer Res. Clin. Oncol.* 138, 647–655.
- [188] Li, J. and Li, G. (2010) Cell cycle regulator ING4 is a suppressor of melanoma angiogenesis that is regulated by the metastasis suppressor BRMS1. *Cancer Res.* 70, 10445–10453.
- [189] Xie, Y.F., Sheng, W., Xiang, J., et al. (2009) Adenovirus-mediated ING4 expression suppresses pancreatic carcinoma cell growth via induction of cell-cycle alteration, apoptosis, and inhibition of tumor angiogenesis. *Cancer Biother. Radiopharm.* 24, 261–269.
- [190] Garkavtsev, I., Kozin, S.V., Chernova, O., et al. (2004) The candidate tumour suppressor protein ING4 regulates brain tumour growth and angiogenesis. *Nature* 428, 328–332.
- [191] Tzouveleakis, A., Ntoliou, P., Karameris, A., et al. (2012) Expression of hypoxia-inducible factor (HIF)-1 $\alpha$ -vascular endothelial growth factor (VEGF)-inhibitory growth factor (ING)-4 axis in sarcoidosis patients. *BMC Res. Notes* 5, 654.
- [192] Zhang, X., Zhu, W., Zhang, J., et al. (2010) MicroRNA-650 targets ING4 to promote gastric cancer tumorigenicity. *Biochem. Biophys. Res. Commun.* 239, 275–280.
- [193] Mraz, M., Dolezalova, D., Plevova, K., et al. (2012) MicroRNA-650 expression is influenced by immunoglobulin gene rearrangement and affects the biology of chronic lymphocytic leukemia. *Blood* 11, 2110–2113.
- [194] Zhang, X.J., Zeng, C.W., Zhang, H., et al. (2010) Dysregulation of miR-15a and miR-214 in human pancreatic cancer. *J. Hematol. Oncol.* 3, 46.
- [195] Huang, J.Y., Cui, S.Y., Chen, Y.T., et al. (2013) MicroRNA-650 was a prognostic factor in human lung adenocarcinoma and confers the docetaxel chemoresistance of lung adenocarcinoma cells via regulating Bcl-2/Bax expression. *PLoS One* 8, e72615.
- [196] Guo, Q. and Fast, W. (2011) Citrullination of inhibitor of growth 4 (ING4) by peptidylarginine deiminase 4 (PAD4) disrupts the interaction between ING4 and p53. *J. Biol. Chem.* 286, 17069–17078.
- [197] Ozer, A. and Bruick, R.K. (2005) Regulation of HIF by prolyl hydroxylases: recruitment of the candidate tumor suppressor protein ING4. *Cell Cycle* 4, 1153–1156.
- [198] Shiseki, M., Nagashima, M., Pedoux, R.M., et al. (2003) p29ING4 and p28ING5 bind to p53 and p300, and enhance p53 activity. *Cancer Res.* 63, 2373–2378.
- [199] Coles, A.H., Gannon, H., Cerny, A., et al. (2010) Inhibitor of growth-4 promotes I $\kappa$ B $\alpha$  promoter activation to suppress NF- $\kappa$ B signaling and innate immunity. *Proc. Natl. Acad. Sci. U.S.A.* 107, 11423–11428.
- [200] Palacios, A., Munoz, I.G., Pantoja-Uceda, D., Marcaida, M.J., et al. (2008) Molecular basis of histone H3K4me3 recognition by ING4. *J. Biol. Chem.* 283, 15956–15964.
- [201] Palacios, A., Moreno, A., Oliveira, B.L., et al. (2010) The dimeric structure and the bivalent recognition of H3K4me3 by the tumor suppressor ING4 suggests a mechanism for enhanced targeting of the HBO1 complex to chromatin. *J. Mol. Biol.* 396, 1117–1127.
- [202] Mishima, Y., Miyagi, S., Saraya, A., et al. (2011) The Hbol-Brdl/Brpf2 complex 1546 is responsible for global acetylation of H3K14 and required for fetal liver erythropoiesis. *Blood* 118, 2443–2445.
- [203] Zhang, F., Baumer, N., Rode, M., et al. (2011) The inhibitor of growth protein 5 (ING5) depends on INCA1 as a co-factor for its antiproliferative effects. *PLoS One* 6, e21505.
- [204] Xing, Y.N., Yang, X., Xu, X.Y., et al. (2011) The altered expression of ING5 protein is involved in gastric carcinogenesis and subsequent progression. *Hum. Pathol.* 42, 25–35.
- [205] Li, X., Kikuchi, K. and Takano, Y. (2011) ING genes work as tumor suppressor genes in the carcinogenesis of head and neck squamous cell carcinoma. *J. Oncol.* 2011, e963614.
- [206] Qi, L. and Zhang, Y. (2013) Truncation of inhibitor of growth family protein 5 effectively induces senescence, but not apoptosis in human tongue squamous cell carcinoma cell line. *Tumour Biol.* [Epub ahead of print].
- [207] Zheng, H.C., Xia, P., Xu, X.Y., et al. (2011) The nuclear to cytoplasmic shift of ING5 protein during colorectal carcinogenesis with their distinct links to pathologic behaviors of carcinomas. *Hum. Pathol.* 42, 424–433.
- [208] Wang, W., Zhu, W., Xu, X.Y., et al. (2012) The clinicopathological significance of REIC expression in colorectal carcinomas. *Histol. Histopathol.* 27, 735–743.
- [209] Liu, M., Du, Y., Gao, J., et al. (2013) Aberrant expression miR-196a is associated with abnormal apoptosis, invasion and proliferation of pancreatic cancer cells. *Pancreas* 42, 1169–1181.
- [210] Wang, J., Huang, W., Wu, Y., et al. (2012) MicroRNA-193 pro-proliferation effects for bone mesenchymal stem cells after low-level laser irradiation treatment through inhibitor of growth family, member 5. *Stem Cells Dev.* 21, 2508–2519.
- [211] Hornbeck, P.V., Kornhauser, J.M., Tkachev, S., et al. (2012) PhosphoSitePlus: a comprehensive resource for investigating the structure and function of experimentally determined post-translational modifications in man and mouse. *Nucleic Acids Res.* 40, D261–D270.
- [212] Liu, N., Wang, J., Wang, J., et al. (2013) ING5 is a Tip60 cofactor that acetylates p53 in response to DNA damage. *Cancer Res.* 73, 3749–3760.
- [213] Mulder, K.W., Wang, X., Escriu, C., et al. (2012) Diverse epigenetic strategies interact to control epidermal differentiation. *Nat. Cell. Biol.* 14, 753–763.
- [214] Champagne, K.S., Saksouk, N., Pena, P.V., et al. (2008) The crystal structure of the ING5 PHD finger in complex with an H3K4me3 histone peptide. *Proteins* 72, 1371–1376.
- [215] Lalonde, M.E., Awakumov, N., Glass, K.C., et al. (2013) Exchange of associated factors directs a switch in HBO1 acetyltransferase histone tail specificity. *Genes Dev.* 27, 2009–2024.
- [216] Kristensen, L.S., Nielsen, H.M. and Hansen, L.L. (2009) Epigenetics and cancer treatment. *Eur. J. Pharmacol.* 625, 131–142.
- [217] Wyce, A., Ganji, G., Smitheman, K.N., et al. (2013) BET inhibition silences expression of MYCN and BCL2 and induces cytotoxicity in neuroblastoma tumor models. *PLoS One* 8, e72967.
- [218] Toyama, T. and Iwase, H. (2004) p33ING1b and estrogen receptor (ER)  $\alpha$ . *Breast Cancer* 11, 33–37.
- [219] Toyama, T., Iwase, H., Yamashita, H., et al. (2003) p33(ING1b) stimulates the transcriptional activity of the estrogen receptor  $\alpha$  via its activation function (AF) 2 domain. *J. Steroid Biochem. Mol. Biol.* 87, 57–63.
- [220] Tamannai, M., Farhangi, S., Truss, M., et al. (2010) The inhibitor of growth 1 (ING1) is involved in trichostatin A-induced apoptosis and caspase 3 signaling in p53-deficient glioblastoma cells. *Oncol. Res.* 18, 469–480.
- [221] Feng, X., Bonni, S. and Riabowol, K. (2006) HSP70 induction by ING1 proteins sensitizes cells to TNF- $\alpha$  receptor mediated apoptosis. *Mol. Cell. Biol.* 26, 9244–9255.
- [222] Takahashi, M., Seki, N., Ozaki, T., et al. (2002) Identification of the p33 (ING1)-regulated genes that include cyclin B1 and proto-oncogene DEK by using cDNA microarray in a mouse mammary epithelial cell line NMuMG. *Cancer Res.* 62, 2203–2209.
- [223] Ishii, N., Maier, D., Merlo, A., et al. (1999) Frequent co-alterations of TP53, p16/CDKN2A, p14ARF, PTEN tumor suppressor genes in human glioma cell lines. *Brain Pathol.* 9, 469–479.
- [224] Tallen, U.G., Truss, M., Kunitz, F., et al. (2008) Down-regulation of the inhibitor of growth 1 (ING1) tumor suppressor sensitizes p53-deficient glioblastoma cells to cisplatin-induced cell death. *J. Neurooncol.* 86, 23–30.
- [225] Zhu, J.J., Li, F.B., Zhu, X.F., et al. (2006) The p33ING1b tumor suppressor cooperates with p53 to induce apoptosis in response to etoposide in human osteosarcoma cells. *Life Sci.* 78, 1469–1477.
- [226] Zhu, J.J., Li, F.B., Zhou, J.M., et al. (2005) The tumor suppressor p33ING1b enhances taxol-induced apoptosis by p53-dependent pathway in human osteosarcoma U2OS cells. *Cancer Biol. Ther.* 4, 39–47.
- [227] Smith, K.T., Martin-Brown, S.A., Florens, L., et al. (2010) Deacetylase inhibitors dissociate the histone-targeting ING2 subunit from the Sin3 complex. *Chem. Biol.* 17, 65–74.

- [228] Zhu, Y., Du, X., Chen, H., et al. (2011) Effect of adenovirus-mediated ING4 and IL-24 co-expression on chemosensitivity to human lung adenocarcinoma in vitro and in vivo. *Sheng Wu Gong Cheng Xue Bao* 27, 85–94.
- [229] Ling, C., Xie, Y., Zhao, D., et al. (2012) Enhanced radiosensitivity of non-small-cell lung cancer (NSCLC) by adenovirus-mediated ING4 gene therapy. *Cancer Gene Ther.* 19, 697–706.
- [230] Mao, Z.L., He, S.B., Sheng, W.H., et al. (2013) Adenovirus-mediated ING4 expression reduces multidrug resistance of human gastric carcinoma cells in vitro and in vivo. *Oncol. Rep.* 30, 2187–2194.
- [231] Wang, R., Huang, J., Feng, B., et al. (2012) Identification of ING4 (inhibitor of growth 4) as a modulator of docetaxel sensitivity in human lung adenocarcinoma. *Mol. Med.* 18, 874–886.
- [232] Xie, Y., Sheng, W., Miao, J., et al. (2011) Enhanced antitumor activity by combining an adenovirus harboring ING4 with cisplatin for hepatocarcinoma cells. *Cancer Gene Ther.* 18, 176–188.
- [233] Mendes-Pereira, A.M., Sims, D., Dexter, T., et al. (2012) Genome-wide functional screen identifies a compendium of genes affecting sensitivity to tamoxifen. *Proc. Natl. Acad. Sci. U.S.A.* 109, 2730–2735.